

Middle to Upper Permian radiolarian faunas from chert blocks in Pai area, northwestern Thailand

NUTTHAWUT WONGANAN¹ & MARTIAL CARIDROIT²

Key words: Radiolaria, chert, biostratigraphy, Permian, northwestern Thailand
Mots clés: Radiolaire, «chert», biostratigraphie, Permien, Thaïlande du Nord-Ouest

ABSTRACT

Well-preserved Permian radiolarians have been discovered from chert blocks embedded in a unit in the Pai district, Mae Hong Son province, northwestern Thailand that was recently mapped as Carboniferous. Twenty-four taxa belonging to ten genera have been recognized and some significant forms of *Follicucullus* are present. Two upper Maokouan (Capitanian) to lower Wuchiapingian radiolarian assemblages (*Follicucullus monacanthus*, *Follicucullus charveti* – *F. porrectus* assemblages) are reported. *Follicucullus charveti* CARIDROIT & DE WEVER is interpreted as a possible provincialism rather than being endemic like other members of the same genus. These results provide more data to demonstrate that in northern Thailand, the Devonian to Triassic was of a zone of deep siliceous marine sedimentation. The resultant deposits are one of the longest witnesses of continuous deposition in an oceanic setting.

RESUME

Des assemblages de radiolaires bien préservés ont été extraits de blocs de «chert» présents dans la région de Pai (Province de Mae Hong Son, Thaïlande du Nord), récemment cartographiés en tant que roches du Carbonifère. Vingt-quatre espèces appartenant à 10 genres sont reconnues, comprenant plusieurs *Follicucullidae* très significatifs pour la stratigraphie. Deux zones d'assemblages (*Follicucullus monacanthus* et *Follicucullus charveti*-*F. porrectus*) sont présentes et datent le Maokouien supérieur (Capitanien) et le Wuchiapingien inférieur; elles sont similaires à celles présentes en Chine du Sud et au Japon. La présence de *Follicucullus charveti* CARIDROIT & DE WEVER, espèce interprétée comme une forme d'eau chaude, est discutée. Ces résultats sont fondamentaux et sont une donnée supplémentaire pour prouver que l'ensemble des temps Dévonien à Trias supérieur est représenté par des dépôts siliceux de mer profonde; ceci étant un des plus longs témoins connus d'une sédimentation océanique continue.

1. Introduction

The study of Permian radiolarians in Thailand, has become a popular topic since radiolarian research started (Caridroit et al., 1990). Occurrences of Permian radiolarians are widespread and they have now been studied in many regions; northern Thailand (Chiang Mai: e.g. Caridroit 1991, 1993; Caridroit et al. 1992; Sashida et al. 1993; Wonganan et al. 2002), north-central Thailand (Sukhothai: Sashida & Nakornsri 1997), north-eastern Thailand (Loei: e.g. Sashida et al. 1993; Sashida & Igo 1999), eastern Thailand (Sra Khaew: e.g. Hada et al. 1999; Sashida et al. 2000). The goals of these studies are both to improve the biostratigraphy and to track radiolarites as remnants of oceanic palaeoenvironments. Only lower to middle Lower and uppermost Permian faunas have been reported from northern Thailand so far (e.g. Caridroit 1993; Sashida et al. 1993). The aim of this paper is to present details of a new upper Middle to lower Upper Permian radiolarian locality

from Thailand. Radiolarian assemblage zones described from Japan and China are identified and briefly discussed. Possible provincialism is observed with the occurrence of *Follicucullus charveti* CARIDROIT & DE WEVER at few localities, compared with the endemic nature of other members of this genus.

2. General Geology and Occurrence of Radiolarians

The district of Pai is located in NW northern Thailand, approximately 110 km from Chiang Mai. According to Bunopas (1981) the main Palaeozoic sedimentary rocks widely exposed in the area is the Mae Hong Son Formation. Age assignments for this unit range from Siluro-Devonian (Bunopas 1981) to Carboniferous-Permian (Chuaviroj et al. 1985; Intawong et al. 1997). The formation is dominated by unmetamorphosed Palaeozoic sedimentary rocks, which are extensively distributed in northwest Thailand. It mainly consists of massive or

¹ Department of Geological Sciences, Faculty of Science, Chiang Mai University, Chiang Mai 50200 Thailand. E-mail: n-thawut@chiangmai.ac.th

² Laboratoire de Paléontologie et Paléogéographie du Paléozoïque (LP3), UMR 8014 et UFR 1818 du C.N.R.S., Sciences de la Terre: Université des Sciences et Technologies de Lille, F-59655 Villeneuve d'Ascq cedex France. E-mail: martial.caridroit@univ-lille1.fr




Fig. 1. Index map of northwestern Thailand showing the studied area and the radiolarian locality.

bedded sandstone, shale, chert and slate (Fig. 2). Limestones, forming isolated karstic hills are widespread and contain abundant Carboniferous and Permian foraminifers and other fossils. In general most Thai geologists correlate these limestones with the Ratburi Group. Limestones in this area are lithologically variable, but mainly consist of massive limestones, with local occurrences of dolomitic limestone with lenticular or nodular cherts. The relationship between limestone units and Palaeozoic siliceous sedimentary units is not clear. Field observations indicate the limestones unconformably overlie siliceous sediments, and where faulted the basal contact is locally marked by breccias.

The studied section is located approximately 7–8 km north of Pai (Fig. 2), or at position 438455 on Thai topographic map sheet 4647 I Amphoe Pai, in the area of Ban Tan Jed Ton village. The section is exposed for approximately 850 m along a gravel road connecting Pai and Wieng Haeng that has been cut through a high mountainous area. The section consists of gray to greenish gray, partly dark gray, well-bedded chert with 4 to 10 cm thick beds intercalated with thin shale and clay (mm – few cm) layers. The main and lowermost exposed section is about 150 m long, before being obscured by strongly weathered yellow volcanic rocks for about 200 m. The second section, which is believed to be a separate block, is composed of greenish gray well-bedded chert and about 10 m long. Ten chert samples were collected from the main section (PAI-391 to PAI-400) and three more were taken from the second (PAI-401 to PAI-403). The next three samples (PAI-404 to PAI-406) were taken from a small chert outcrop exposed on the other side of 200 m of weathered tuffaceous material. Four chert samples were collected (PAI-407 to PAI-412) from the last section which ends at a summit approximately 150 m from the previous section. This chert has long been regarded as chert

beds intercalated within the Upper Silurian to Carboniferous Mae Hong Son Formation (Bunopas 1981), or newly established Carboniferous-Permian of Chauviroj et al. (1985). According to field observations, no outcrops were found which expose the strata underlying the chert.

Under the microscope, the chert from lower part consists of mainly cryptocrystalline to microcrystalline quartz associated with very fine clay minerals. Radiolarians are rather well preserved and abundant and filled by chalcedony. Further up the lithologic column the cherts consist of cryptocrystalline quartz with more abundant fine clay mineral and fine volcanic grains, without any traces of fossils. This probably indicates that there was volcanic activity within or nearby the site of deposition area conditions did not favour the preservation of radiolarian skeletons.

Samples from the main chert section (PAI-391 through PAI-400) yielded the following Upper Permian radiolarian faunas: *Follicucullus ventricosus* ORMISTON & BABCOCK, *F. scholasticus* ORMISTON & BABCOCK, *F. orthogonus* CARIDROIT & DE WEVER, *F. porrectus* RUDENKO, *F. charveti* CARIDROIT & DE WEVER, *F. sp. cf. F. bipartitus* CARIDROIT & DE WEVER, *Hegleria mammilla* (SHENG & WANG) and others. The association of *F. charveti* and *F. porrectus* RUDENKO is diagnostic for assignment to the lower Wuchiapingian. Sample PAI-403 contains fewer radiolarians, however the occurrence of *F. monacanthus* ISHIGA & IMOTO is the diagnostic of the Middle Permian *F. monacanthus* Assemblage (Capitanian), therefore these chert beds are assigned to the Capitanian.

3. Radiolarian biostratigraphy

Twenty-two chert samples were collected, and nine samples yielded Permian radiolarians (Plate 1). The studied chert sec-




Fig. 2. Simplified geologic map of Pai area, Mae Hong Son province, northwestern Thailand (see Fig. 1 for location); shows the studied sections. (after Geologic map 1:50,000 Sheet Amphoe Pai Quadrangle, Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand 1985; Intawong et al. 1997, modified)

tion extends over two radiolarian assemblage zones: the *Follicucullus monacanthus*, and *Follicucullus charveti* – *F. porreectus* assemblage zones. These zones have been reported by Ishiga (1986, 1990) from Japan, Wang et al. (1994) from China and Caridroit in De Wever et al. (2001) and are known to occur in the middle Maokouan (Capitanian) to lower Wuchiapingian.

Follicucullus monacanthus assemblage zone.

This assemblage zone is found within sample PAI-403. It is characterized by the occurrence of *Follicucullus monacanthus* ISHIGA & IMOTO, which is the characteristic taxon. Other radiolarians species that occur in this assemblage include *F. scholasticus* ORMISTON & BABCOCK, *Pseudotormentus* sp. cf. *P. kamigoriensis* CARIDROIT & DE WEVER, *F. orthogonus* CARIDROIT & DE WEVER, *F. ventricosus* ORMISTON & BABCOCK, and *Ormistonella robusta* CARIDROIT & DE WEVER. The *Follicucullus monacanthus* assemblage is known from Japan (e.g. Ishiga 1986), Far East Russia (Rudenko & Panasenko

1990), Oregon in North America (Blome & Reed 1992), South China (Wang et al. 1994), and is also reported in studies by Caridroit (in De Wever et al. 2001, fig. 202, p. 313).

Follicucullus charveti – *F. porreectus* assemblage zone

This assemblage zone is found within green chert samples PAI-393, 396, 398 & 399. It is defined by the co-occurrence of *Follicucullus charveti* CARIDROIT & DE WEVER and *F. porreectus* RUDENKO. The range of this assemblage is considered as a total range of *F. charveti* CARIDROIT & DE WEVER. The other characteristic species are *Follicucullus ventricosus* ORMISTON & BABCOCK, *Follicucullus scholasticus* ORMISTON & BABCOCK, *Follicucullus* sp. cf. *F. bipatitus* CARIDROIT & DE WEVER, *Ishigaum similicutis* CARIDROIT & DE WEVER, *Triplanospongia musashiensis* SASHIDA & TONISHI, *Latentifistula texana* NAZAROV & ORMISTON, *Hegleria mammilla* (SHENG & WANG), and others. The assemblage contains the same radiolarian species reported from Japan (e.g. Ishiga 1990), North




Fig. 3. Columnar sections of the studied bedded chert section exposed on road between Pai and Wieng Hang cities in Ban Tan Jed Ton area (Pai city, Mae Hong Son province), showing distribution of characteristic taxa of radiolarians obtained.

America (e.g. Murchey 1990), South China (e.g. Wang et al. 1994) and New Zealand (e.g. Caridroit & Ferriere 1988). It is comparable to the *F. scholasticus* – *F. venticosus* zone from China (Wang et al. 1994) and the *F. venticosus* – *Ps. fusiformis* assemblage of North America (Murchey 1990).

4. Discussion: *Follicucullus* faunas and palaeobiogeography

Follicucullus charveti CARIDROIT & DE WEVER is characterized by a long apical cone, which is slightly curved to the ventral side (Pl. 1; Fig. 8). Specimens from the area we studied have a strongly inflated pseudothorax and a short pseudoabdomen with small, disk-like, rather short ventral flap compared to those reported from Japan and S. China (e.g. Caridroit & De Wever 1986; Wang & Li 1994). *F. scholasticus* ORMISTON & BABCOCK m. I (e.g. Ishiga 1985; Blome & Reed 1992; Wang et al. 1994), and *F. scholasticus* ORMISTON & BABCOCK m. II (e.g. Ishiga 1985) are considered as two different species. The first is the most abundant form and found in all levels of the section. This form has a smooth, undulating shell, however, somewhat obscure separated portions can be observed in some specimens. Small very short flaps, extending at apertural margin of the dorsal and ventral sides, are well preserved on most specimens. The second morphotype, on the other hand, has a bulbous shell characteristic similar to those reported from Japan

by Caridroit & De Wever (1986) and from the Prymorye region, Far East Russia by Rudenko & Panasenko (1990) and is here regarded as *F. porrectus* RUDENKO. Specimens of *Follicucullus venticosus* ORMISTON & BABCOCK (Pl. 1; Fig. 9) from the study area are closely similar to those reported from N. America (Ormiston & Babcock 1979; Blome & Reed 1992) and from S. China (Wang et al. 1994) having a slightly curved apical cone and big pseudothorax, with short pseudoabdomen; both ventral and dorsal flap are very short.

Materials presented in this study also contains some broken pieces that have been assigned to species such as *Gustefana obliqueannulata* KOZUR (Pl. 1; Fig. 30) and *Nazarovella phlogidea* WANG (Pl. 1; Fig. 31). These forms in fact have little taxonomic value and in our collections all of these broken pieces appear to be the ending part of an arm of already defined radiolarian species.

It is not yet proven that siliceous skeleton zooplanktonic faunas are useful indicators for palaeo-ecology and/or palaeogeographic position. Nevertheless, the possible palaeogeographic implications of the occurrence of a particular taxon with distinctive form at a few localities merit some discussion. The Upper Permian *Follicucullus charveti* – *F. porrectus* radiolarian assemblage recognized in this study is believed to be equivalent to the *F. bipartitus*- *F. charveti* assemblage known from southwest Japan (e.g. Caridroit & De Wever 1984, 1986),




Fig. 4. Palaeogeographic reconstruction during Middle Permian (late Capitanian) showing relative palaeogeographic positions and possible *Follicucillus charveti* province discussed in the text. (Palaeogeographic map and terrane positions are based on Scotese 1997; Metcalfe 1996, and water province areas are after Henderson & Mei 2003, where ST: Shan-Thai; T: Tarim; WB: West Burma).

northern New Zealand (e.g. Caridroit & Ferrière 1988; Take-mura et al. 1999), and south & southwest China (e.g. Wang & Li 1994). According to the known literature reported so far, this assemblage has only been discovered from the four different geographic areas mentioned above. Regarding *Follicucillus charveti* CARIDROIT & DE WEVER, a diagnostic taxon of this assemblage, it is not clear whether it occurs in a restricted area with short period (early Wuchiapigian). The occurrence of this fauna in a restricted area as plotted on a palaeogeographic map may suggest that its geographic range was probably confined to a special part in the tropical palaeoequator (low-latitude) realm within the equatorial warm water province. As this species can likely be defined as a warm-water taxon the palaeogeographic positions of terranes containing radiolarites with this fauna were probably at low latitudes not far from one another (Fig. 4).

Acknowledgements

Our field survey in January 2003 was financial supported by the Chiang Mai University (Thailand), the Laboratoire de Paléontologie et Paléogéographie du Paléozoïque (CNRS-UMR 8014) and the Université des Sciences et Technologies de Lille (France). The Faculty of Science, Chiang Mai University for supporting field-working facilities is credited here. Fabrice Corday and Frances Spiller are thanked for reviews that helped to improve the manuscript.

REFERENCES

BLOME, C. D. & REED, K. M. 1992: Permian and Early (?) Triassic radiolarian faunas from the Grindstone terrane, central Oregon. *Journal of Paleontology* 66, 351–383.

BUNOPAS, S. 1981: Palaeogeographic history of western Thailand and adjacent parts of South-east Asia: a plate tectonic interpretation. Geological Survey paper vol. 5, Department of Mineral Resources, Bangkok, Special Issue, 810.

CARIDROIT, M. 1991: Taxonomic study on Carboniferous and Permian Radiolaria from NW Thailand. Paleontologic, stratigraphic and tectonic significances. Abstracts. Sixth Meeting, International Association of Radiolarian Paleontologists (INTERRAD VI), p. 21.

CARIDROIT, M. 1993: Permian Radiolarian from NW Thailand. In: THANASUTHIPITAK, T. (Ed.): Proceedings of the International Symposium on Biostratigraphy of Mainland Southeast Asia Facies and Paleontology 1, Chiang Mai, Thailand, 83–96.

CARIDROIT, M. & DE WEVER, P. 1984: Description de quelques nouvelles espèces de Follicucullidae et d'Entactinidae (Radiolaires polycystines) du Permien du Japon (Description of some new species of Follicucullidae and Entactinidae (Polycystin radiolarians) from the Permian of Japan). *Geobios* 17, 639–644.

CARIDROIT, M. & DE WEVER, P. 1986: Some Late Permian Radiolarians from pelitic rocks of the Tatsuno Formation (Hyogo Prefecture), Southwest Japan. *Marine Micropaleontology* 11, 55–90.

CARIDROIT, M. & FERRIERE, J. 1988: Premières datations précises du Paléozoïque par radiolaires en Nouvelle Zélande: intérêt géologique et paléontologique. *Compte rendus Académie des Sciences de l'Académie des Sciences (Paris), series II* 306, 321–326.

CARIDROIT, M., FONTAINE, H., JONGKANJAMASOONTORN, Y., SUTEETHORN, V. & VACHARD, D. 1990: First results of a paleontological study of Northwest Thailand. *CCOP Technical Secretariat*, 337–350.

CARIDROIT, M., VACHARD, D. & FONTAINE, H. 1992: Datations par radiolaires (Carbonifère, Permien et Trias) en Thaïlande nord-occidentale. Mise en évidence de nappe de charriage et d'olisostromes. *Académie des Sciences Compte Rendu Paris* 314, 515–520.

CHAUVIROJ, S., CHATURONGKAWANICH, S., LEEVONGCHAROEN, S. & SOPONPONGPIPAT, P. 1985: Hydrothermal Research Project, Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand. 66.

DE WEVER, P., DUMITRICA, P., CAULET, J.P., NIGRINI, C. & CARIDROIT, M. 2001: Radiolarians in the Sedimentary Record. 533 p., Gordon and Breach Science Publishers.

HADA, S., BUNOPAS, S., ISHII, K. & YOSHIKURA, S. 1999 Rift-drift history and the amalgamation of Shan-Thai and Indochina/East Malaya blocks In: METCALFE, I. (Ed.): Gondwana Dispersion and Asian Accretion. *IGCP 321, Sp. Publ.*, 67–88, A.A. Balkema.

HENDERSON, C.M. & MEI, S. 1990: Stratigraphic versus environment significance of Permian serrated conodonts around the Cisuralian-Guadalupian boundary: new evidence from Oman. *Palaeogeography, Palaeoclimatology, Palaeoecology* 191, 301–328.

INTAWONG, A., KUNMANEE, J. & WONGANAN, N. 1997: Geological report in the area of Ban Tan Jed Ton, Tambol Wieng Nue, Amphoe Pai, Chang Wat Mae Hong Son. Department of Geological Sciences, Chiang Mai University, 82.

ISHIGA, H. 1985: Discovery of Permian radiolarians from Katsumi and Oi Formation along south of Maizuru Belt, Southwest Japan and its significance. *Ibid* 39, 175–185.

- ISHIGA, H. 1986: Late Carboniferous and Permian radiolarian biostratigraphy of southwest Japan. *Journal of Geoscience*, Osaka City University 29, 89–100.
- ISHIGA, H. 1990: Paleozoic radiolarians. In: ICHIKAWA, K., MIZUTANI, S., HARA, I., HADA, S. & YAO, A. (Eds.): Pre-Cretaceous terranes of Japan. Publication of IGCP Project No. 224: Pre-Jurassic Evolution of Eastern Asia, Nippon Insatsu Shuppan, Co. Ltd, Osaka, 285–295.
- METCALFE, I. 1996: Gondwana dispersion, Asian accretion and evolution of Eastern Tethys. *Australian Journal of Earth Sciences* 43, 605–623.
- MURCHEY, B.L. 1990: Age and Depositional setting of siliceous sediments in the Upper Paleozoic Havallah sequence near Battle Mountain, Nevada: implications for the paleogeography and structural evolution of the western margin of North America. *Geological Society of America, Special paper* 255, 137–155.
- ORMISTON, A. & BABCOCK, L. 1979: *Follicucullus*, new radiolarian genus from the Gaudalupian (Permian) Lamar Limestone of the Delaware Basin. *Journal of Paleontology* 53, 328–334.
- RUDENKO, V. & PANASENKO, E.S. 1990: A new findings of the Upper Permian radiolarians in Prymorye region. In: ZAKHAROV, Y.D., BELYAEVA, G.V. & NIKITINA, A.P. (Eds.): New data on Paleozoic and Mesozoic biostratigraphy of the South Far East. Far Eastern Branch of the USSR Acad. of Sci., Vladivostok, 117–124.
- SASHIDA, K. & IGO, H. 1999: Occurrence and tectonic significance of Paleozoic and Mesozoic Radiolaria in Thailand and Malaysia. In: METCALFE, I. (Ed.): Gondwana Dispersion and Asian Accretion Final Results Volume for IGCP Project 321, A.A. Balkema, Rotterdam, 175–196.
- SASHIDA, K. & NAKORNSRI, N. 1997: Lower Permian radiolarian faunas from the Khanu Chert Formation distributed in the Sukhothai Area, northern central Thailand. In DHEERADILOK, P., C. HINTHONG, P. CHAODUMRONG, P. PUTTHAPIBAN, W. TANASTHIEN, C. UTHA-AROON, N. SATTYARAK, N. NUCHANONG & T. TECHAWAN (Eds.): Proceedings of the International Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific, Bangkok, Thailand, 101–108.
- SASHIDA, K., IGO, H., HISADA, K., NAKORNSRI, N. & AMPORNMAHA, A. 1993: Occurrence of Paleozoic and early Mesozoic Radiolaria in Thailand (preliminary report). *Journal of Southeast Asian Earth Science* 8, 97–108.
- SASHIDA, K., IGO, H., UENO, K., NAKORNSRI, N. & SARDSUD, A. 1998: Late Paleozoic radiolarian fauna from northern and northeastern Thailand. *Science Reports of the Institute of Geoscience, University of Tsukuba, Section B* 19, 1–27.
- SASHIDA, K., IGO, H., ADASHI, S., UENO, K., KAJIWARA, Y., NAKORNSRI, N. & SARDSUD, A. 2000: Late Permian and Middle Triassic radiolarian faunas from northern Thailand. *Journal of Paleontology* 74, 789–811.
- SCOTSE, K. 1997: Paleogeographic Atlas Paleomap Progress Report 90-0497. Paleomap Project ed., Univ. of Texas at Arlington, Arlington, Texas, 45.
- TAKEMURA, A., MORIMOTO, T., AITA, Y., HORI, R.S., HIGUCHI, Y., SPORLI, K.B., CAMPBELL, H.L., KODAMA, K. & SAKAI, T. 1999: Permian *Albaillellaria* (Radiolaria) from a limestone lens at the Arrow Rocks in the Waipapa Terrane, Northland, New Zealand. In: DE WEVER, P. & CAULET, J.-P. (Eds.): *InterRad VII*, Paris/Bierville, September 1997. *Geodiversitats* 21(4), 789–811.
- WANG, Y. & LI, J. 1994: Discovery of the *Follicucullus bipartitus* – *F. charveti* radiolarian assemblage zone and its geological significance. *Acta Micropalaeontologica Sinica* 11, 201–212.
- WANG, Y., CHENG, Y.N. & YANG, Q. 1994: Biostratigraphy and systematics of Permian radiolarians in China. *Paleoworld* 4, 172–202.
- WONGANAN, N., CARIDROIT, M. & RANDON, C. 2002: Radiolarians and conodonts from radiolarites in NW-Thailand; witnesses of a 140 my (at least) oceanic realm. In: The 64th Palaeontological Association Annual Meeting, Cambridge, abstract. *The Palaeontology Newsletter* 51, 73.

Manuscript received January 2004

Revision accepted March 2005

Plate 1

Permian radiolarians from Pai area, northwestern Thailand. All figures are scanning electronic micrographs. Scale bar 100 µm applies to all specimens. 1–6. *Follicucullus scholasticus* ORMISTON & BABCOCK, PAI-421; 7. *Follicucullus porrectus* RUDENKO, PAI-421; 8–10. *Follicucullus charveti* CARIDROIT & DE WEVER, PAI-421; 11. *Follicucullus monacanthus* ISHIGA & IMOTO, PAI-421; 12–13. *Follicucullus ventricosus* ORMISTON & BABCOCK, PAI-419; 14. *Follicucullus orthogonus* CARIDROIT & DE WEVER, PAI-421; 15–17. *Pseudoalbaillella* sp. A, PAI-421; 18. *Neobaillella* sp. cf. *N. grypa* ISHIGA ET AL., PAI-421; 19. *Albaillella* sp. aff. *A. levis* ISHIGA ET AL., PAI-421; 20. *Nazarovella gracilis* DE WEVER & CARIDROIT, PAI-402; 21. *Quadricaulis femoris* CARIDROIT & DE WEVER, PAI-393; 22. *Ishigaum* (?) *similicutis* CARIDROIT & DE WEVER, PAI-393; 23. *Ishigaum trifustus* DE WEVER & CARIDROIT, PAI-393; 24. *Trianospongus musashiensis* SASHIDA & TONISHI, PAI-393; 25. *Latentifistula texana* NAZAROV & ORMISTON, PAI-398; 26. *Ormistonella robusta* DE WEVER & CARIDROIT, PAI-393; 27. *Pseudotormetus* sp. cf. *P. kamigoriensis* DE WEVER & CARIDROIT, PAI-402; 28. *Copicyntra* sp., PAI-393; 29. *Hegleria mamma* (SHENG & WANG), PAI-393; 30–31. broken part of latentifistulinid (30: cf. *Gustefana obliqueannulata* KOZUR; 31: cf. *Nazarovella phlogidea* WANG), PAI-393.

