

# Ivrea mantle wedge, arc of the Western Alps, and kinematic evolution of the Alps–Apennines orogenic system

Stefan M. Schmid<sup>1</sup>  · Eduard Kissling<sup>1</sup> · Tobias Diehl<sup>1</sup> · Douwe J. J. van Hinsbergen<sup>2</sup> · Giancarlo Molli<sup>3</sup>

Received: 6 June 2016 / Accepted: 9 December 2016 / Published online: 29 March 2017  
© Swiss Geological Society 2017

**Abstract** The construction of five crustal-scale profiles across the Western Alps and the Ivrea mantle wedge integrates up-to-date geological and geophysical information and reveals important along strike changes in the overall structure of the crust of the Western Alpine arc. Tectonic analysis of the profiles, together with a review of the existing literature allows for proposing the following multistage evolution of the arc of the Western Alps: (1) exhumation of the mantle beneath the Ivrea Zone to shallow crustal depths during Mesozoic is a prerequisite for the formation of a strong Ivrea mantle wedge whose strength exceeds that of surrounding mostly quartz-bearing units, and consequently allows for indentation of the Ivrea mantle wedge and eastward back-thrusting of the western Alps during Alpine orogeny. (2) A first early stage (pre-35 Ma) of the West-Alpine orogenic evolution is characterized by top-NNW thrusting in sinistral transpression causing at least some 260 km displacement of internal Western Alps and E-W-striking Alps farther east, together with the Adria micro-plate, towards N to NNW with respect to stable Europe. (3) The second stage (35–25 Ma), further accentuating the arc, is associated with top-WNW thrusting in the external zones of the central portion of the arc and is

related to the lateral indentation of the Ivrea mantle slice towards WNW by some 100–150 km. (4) The final stage of arc formation (25–0 Ma) is associated with orogeny in the Apennines leading to oroclinal bending in the southernmost Western Alps in connection with the 50° counter-clockwise rotation of the Corsica-Sardinia block and the Ligurian Alps. Analysis of existing literature data on the Alps–Apennines transition zone reveals that substantial parts of the Northern Apennines formerly suffered Alpine-type shortening associated with an E-dipping Alpine subduction zone and were backthrusted to the NE during Apenninic orogeny that commences in the Oligocene.

**Keywords** Western Alps · Apennines · Ivrea mantle · Teleseismic tomography · Arcuate mountain belts

## 1 Introduction

The Western Alps represent a prime example of a very arcuate mountain belt curving around from an E-W strike with dominant top-N nappe stacking in the Eastern Alps to a N-S strike in the northern Western Alps and finally into an E-W strike with top-S nappe stacking in the Ligurian Alps straddling the Mediterranean coast (Fig. 1). The question after the mechanisms of arc formation is an old and much debated issue in many orogens (Marshak 1988) involving, in the case of the Alps, elements of oroclinal bending of an initially straight orogen (Carey 1955; van der Voo 2004), induced by rigid body rotation of the hinterland (Vialon et al. 1989) and/or by W-directed indentation of the Adria microplate (Laubscher 1971; Schmid and Kissling 2000). Transpressional movements associated with large top NNW displacements of portions of the inner zone of the Western Alps that occur at a small angle to the present-day N-S strike


Editorial handling: C. Sue and A. G. Milnes.

✉ Stefan M. Schmid  
stefan.schmid@unibas.ch

<sup>1</sup> Institute of Geophysics, ETH Zürich, Sonneggstrasse 5, 8092 Zurich, Switzerland

<sup>2</sup> Department of Earth Sciences, Utrecht University, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands

<sup>3</sup> Dipartimento di Scienze della Terra, Università di Pisa, Pisa, Italy



**Fig. 1** Tectonic map of the arc of the Western Alps after Schmid et al. (2004) and Bousquet et al. (2012b) with indication of the traces of the profiles depicted in Fig. 2. The selected faults indicated in red are: RSL Rhone-Simplon Line, PL Periadriatic Line, PF Penninic

Front, ACL Argentera-Cuneo Line, AFSV front of the metamorphic core of the Alps and Sestri-Voltaggio line, VVL Villalvernia-Varzi line. Abbreviations in black are: LA Ligurian Alps, TH Torino hills, MH Monferrato hills, GB Giudicarie transpressive belt

of the Western Alps were also invoked for being responsible for explaining the curvature from the Eastern Alps into the N-S striking Western Alps (Ricou and Siddans 1986). There have also been attempts to explain diverging kinematic indicators along the arc of the Western Alps ranging from top-N Western Switzerland to top-SW in southern France without taking resort to a multi-stage evolution allowing for strike-slip movements and/or large rotations (Platt et al. 1989; their group 3 kinematic indicators formed in the 40–15 Ma time interval). However, as pointed out by Butler (1986; his Figure 2), strongly divergent thrust transport directions over large distances inevitably result in excessive orogen-parallel stretching.

In the Western Alps many previous workers (e.g. Laubscher 1971; Butler 1983) recognized the crucial role of the Ivrea mantle wedge residing at relatively shallow crustal levels and locally reaching the surface. Novel geophysical methods such as teleseismic tomography aiming at the structure of the earth's mantle (Lippitsch et al. 2003) and crustal tomography based on local earthquake data (Diehl et al. 2009), together with older information based on controlled source seismology (Blundell et al. 1992), led to

substantial recent improvements in revealing the deep structure of the arc of the Western Alps. The new geophysical data set of unprecedented quality calls for integration with a wealth of geological data collected over many decades to elucidate the formation of the arc of the Western Alps that also demands to analyze the Alps–Apennines transition area located in the Ligurian Alps and the Northern Apennines (Molli et al. 2010).

This contribution integrates the new results of crustal tomography and improved knowledge on Moho depth (Spada et al. 2013) with a geological interpretation of five crustal-scale transects across the Western Alps. Thereby we focus on the role of the Ivrea mantle wedge. Another important aim of our geological-geophysical interpretations is to put new constraints on the question as to how much continental crust was subducted in the Alps and what the variations in pre-orogenic crustal thickness might have been (see discussions by, e.g., Ménéard and Molnar 1991; Butler 2013; Mohn et al. 2014).

Two transects are revised versions of previously published profiles (Schmid et al. 2004) while three of them were newly constructed. Special emphasis is given to

improved mapping of the outlines of the Ivrea mantle wedge based on geophysical data. This is then followed by an analysis of the complex multi-stage evolution of the arc of the Western Alps and an attempt to map the associated displacement history of geological units of the Western Alps–Northern Apennines system using plate reconstruction methodology (GPlates free software of Boyden et al. 2011) embedded in the global plate kinematic reconstruction of Seton et al. 2012. Finally, we provide a review-type discussion of the Alps–Apennines transition area and discuss controversial issues concerning questions of multi-stage vs. single subduction systems in the Apennines.

## 2 Crustal-scale profiles across the Western Alps and the Ivrea mantle wedge integrating geological and geophysical information

To constrain the geometry of the Ivrea mantle wedge, and as a basis for discussing its role during the formation of the arc of the Western Alps, we constructed a series of crustal scale profiles along the arc of the Western Alps (Fig. 1). These profiles integrate geological data with recent geophysical data focusing on the deep structure. The construction of the profiles is based on a series of assumptions, the most important one being that advocated by Schmid and Kissling (2000), namely that the European lower crust is decoupled from the upper crust and goes into subduction together with the underlying mantle for considerable distance below the Ivrea mantle wedge. This invokes a basal detachment of the external thrust sheets at the interface between upper and lower crust at a depth of 20–30 km. While this assumption is compatible with the velocity structure (see below) and recent geophysical findings of Zhao et al. (2015) it is by no means generally accepted (see discussion in Butler 2013). However, there are considerable uncertainties about the exact pre-orogenic thickness of the European and Briançonnais continental crust in more internal parts of the profiles and this is one reason why we restrain from area balancing over the entire length of our transects. The main reason is that transport directions changed with time during the formation of the arc of the Western Alps, making 2D area balance along the transects impossible.


The geological literature used for constructing the individual profiles varies and hence will be cited when presenting the profiles. However, we used one and the same set of geophysical data for constraining the deep structure in all the profiles. Firstly, we used the results of a tomographic high-resolution 3-D P-wave model of the Alpine crust obtained by Diehl et al. (2009). This 3-D velocity model was used for constructing parts of the sections of Fig. 2 that present contours of equal P-wave velocity superimposed with a geological interpretation.

When comparing the velocity contours with the geological interpretation it has to be kept in mind, however, that these contours have a low spatial resolution since they are based on assigning a given velocity to a sizable cell volume of  $20 \times 20$  km (horizontal) and 15 km (vertical). Secondly, we complemented these tomographic data by geophysical data better constraining Moho-depth. Where available we took Moho depths from Spada et al. (2013; their Figure 11; see also Wagner et al. 2012 regarding the methodology) who combined results of controlled-source seismology with local receiver functions Moho depth maps and local earthquake tomography-derived Moho maps.

Comparison between Moho depth compiled by Spada et al. (2013), characterized by a P-wave velocity jump, and the velocity model of Diehl et al. (2009) that defines Moho depth in terms a high velocity gradient in the range of  $6.5\text{--}8.0$  km  $\text{s}^{-1}$ , noticeable over large parts of the profiles, shows that on average the Moho as defined by Spada et al. (2013) best corresponds with the velocity contour of  $7.25$  km  $\text{s}^{-1}$  in areas characterized by a subhorizontal or only weakly inclined Moho (Diehl et al. 2009). In order to define the strongly inclined boundaries of the Ivrea mantle wedge at shallower depth  $<30$  km, not well constrained by the compilation of Spada et al. (2013), we followed the  $7.0$  km  $\text{s}^{-1}$  contour in all profiles.

In the profiles ECORS-CROP, NFP20-West and Ticino rather thin slices of the Ivrea wedge and/or the lower crust of the Adria microplate, geologically speaking the Ivrea Zone (Handy and Zingg 1991), reach the surface. In these profiles the outlines of the Ivrea mantle wedge are too thin to be resolved by the P-wave velocity model of Diehl et al. (2009) given its limited spatial resolution. The geometry of the very thin and strongly dismembered Ivrea mantle wedge and Adriatic lower crust is only tentatively drawn, being only constrained by the surface exposures.

Another problem one faces when determining the shape of the Ivrea mantle wedge is that the velocities  $>7.25$  km  $\text{s}^{-1}$  expected for the Ivrea mantle rocks that are part of the Adriatic microplate and the adjacent European mantle become indistinguishable from velocities expected for potential remnants of the subduction channel separating European and Ivrea (=Adriatic) mantle marking the present-day plate boundary at great depth. At a depth of 30–45 km and temperatures around  $400\text{--}500$  °C, we expect to reach blueschist facies conditions, the exact depth depending on temperature (Oberhänsli 2004). Based on thermal modeling (Bousquet et al. 1997) the expectation of temperatures between  $450^\circ$  and  $500$  °C at a depth of 30 km is realistic. The P-wave velocity of blueschist rocks at  $400$  °C and 1 GPa is around  $7.25$  km  $\text{s}^{-1}$  on average according to Fujimoto et al. (2010). From a depth of around 50 km onward, again depending on temperature, we expect eclogite facies conditions associated with even higher





◀ **Fig. 2** Five selected geological-tectonic transects across the Western Alps. The sections are superimposed with and partly based on vertical cross sections across the final P-wave velocity model of Diehl et al. (2009) indicated by *red dashed lines*. The Moho (*blue line*) is taken from a combination of controlled source seismology, local earthquake tomography and receiver function analysis (Wagner et al. 2012; Spada et al. 2013). See text for data sources and further discussion and Fig. 1 for the traces of the profiles. **a** Argentera transect; **b** Pelvoux transect; **c** ECORS-CROP transect; **d** NFP 20-West transect; **e** Ticino transect

velocities around  $8.0\text{--}8.5\text{ km s}^{-1}$  (Worthington et al. 2013). Hence, the contours of velocities  $>7.25\text{ km s}^{-1}$  are expected to cross over from the European to the Ivrea mantle rocks traversing the subduction channel formed by high-P metamorphic rocks. Such crossover is actually seen in all the profiles of Fig. 2, also for velocity contours at 7.0, 6.5 and occasionally at  $6.5\text{ km s}^{-1}$ . This is because velocities expected for the subducted European lower crust (around  $6.5\text{ km s}^{-1}$ ; Ye et al. 1995) are similar to those expected for the stack of nappes that is northerly adjacent to the Ivrea mantle wedge, nappes that are only partially overprinted by high-P metamorphism (e.g. Henry et al. 1993). All this, together with the limited resolution of the 3-D velocity model, puts severe limits to a straightforward correlation of velocities with geological-tectonic entities, particularly at greater depths.

## 2.1 Argentera profile (Fig. 2a)

The external SW part of this profile is characterized by minor shortening of around 21 km in Mesozoic sediments (Chaînes Subalpines) of the European foreland (Lickorish and Ford 1998), associated with rather modest late-stage up-thrusting of the Argentera external massif (Malaroda et al. 1970) during the Neogene. In this external part of the profile the European Moho coincides with a strong velocity gradient between 6.5 and  $7.5\text{ s}^{-1}$  in the velocity model of Diehl et al. (2009).

The still unfolded Chaînes Subalpines were thrust by sedimentary units attributed to the Subbriançonnais, together with non-metamorphic Helminthoid flysch detached from the Piedmont-Liguria paleogeographical domain above ophiolite-bearing mélanges (Autapie and Parpaillon nappes; Merle and Brun 1984). Final emplacement of these higher nappes occurred during the Late Eocene (Kerckhove 1969), i.e. before thrusting along the late Alpine (Early Oligocene) out-of-sequence thrust called “Penninic Front” and Neogene shortening in the Chaînes Subalpines. Since the Autapie and Parpaillon nappes lack metamorphic overprint they are considered as units derived from a part of the Piedmont-Liguria Ocean that always remained in an upper plate position. They form relics of the front of the overriding oceanic upper plate (Piedmont-Liguria Upper Plate units of Fig. 2a) that started to thrust the oceanic

lower plate (Piedmont-Liguria Lower Plate units of Fig. 2a) from the late Cretaceous onwards, reaching the front of the Alps at the end of the Eocene, together with the underlying Subbriançonnais units.

There is a rather abrupt lateral change of velocities at a given depth across the boundary between external and internal Alps. This change coincides with the downward projection of the southern end of a very steeply NNE dipping late Alpine out-of-sequence thrust well documented in the Pelvoux section and known as the Penninic Front (Ceriani et al. 2001; Lardeaux et al. 2006; Loprieno et al. 2011; see description of the Pelvoux section). The term “Penninic Front”, synonymous with the term “Briançonnais Frontal Thrust” coined by Tricart (1986) who first recognized the importance of this Early Oligocene out-of-sequence thrust in marking the boundary between internal and external parts of the Western Alps, is a misnomer in that earlier emplaced Penninic units are occasionally also found in front of this out-of-sequence thrust (Ubaye-Embrunais of profile in Fig. 2; Préalpes Romandes of Fig. 2d; see also Fig. 1). We use the term Penninic Front nevertheless because it became deeply entrenched in the literature by now. This top WMW Penninic Frontal Thrust turns into the sinistral Acceglio-Cuneo Line (Fig. 1) and is characterized by a sinistrally transpressive component as it swings around into the WNW-ESE strike of the Argentera-Cuneo line whose western end is traversed by the Argentera profile. The sinistral Acceglio-Cuneo Line allows for the WNW-directed indentation of the Ivrea mantle wedge along its southern boundary located near Cuneo (Trullenque 2005; Ceriani et al. 2001, see their Figure 1). Laubscher (1971) first proposed the existence of a sinistral strike slip zone (his “Ligurian shear zone”) allowing for westward indentation of the Ivrea mantle wedge for kinematic reasons (see also Laubscher 1991). Based on the interpretation of fission track data (Fügenshuh and Schmid 2003) and radiometric dating of mylonites (Simon-Labric et al. 2009) the thrust marking the Penninic Front (Roselend thrust of Loprieno et al. 2011) was active during the Early Oligocene (34–30 Ma). By downward extrapolation the steep dip of this crustal scale transpressive fault projects (Malaroda et al. 1970) into parallelism with the steeply overturned  $6.5\text{ s}^{-1}$  velocity contour that marks the SW edge of a high-velocity area with  $>6.5\text{ s}^{-1}$  located within the high-pressure Briançonnais units at a depth of  $>10\text{ km}$ . This high velocity area is interpreted to consist of the high-pressure Briançonnais units that include the Dora Maira “massif” exposed at the surface at the eastern margin of the Internal Alps (Henry et al. 1993). We are aware that not all exposed parts of the Dora Maira massif constituting the high pressure Briançonnais show blueschist facies or eclogitic parageneses (Bousquet et al. 2012a). However, the high velocities




Fig. 2 continued

at depth require that most of the volume taken up by the Briançonnais at depth must have undergone high-pressure metamorphism.

The shallow crustal parts of the metamorphic core of the Alps in this profile are characterized by large-scale back-folds that formed between 35 and 30 Ma (Bucher et al. 2004) and which affected pre-existing nappe contacts

between: (1) the high-pressure Briançonnais units, (2) overlying so-called Aceglgio and/or Pre-Piémontais units (Lemoine and Tricart 1986; Caby 1996), representing the former transition of the Briançonnais micro-continent into the Piedmont-Liguria Ocean, (3) Piedmont-Liguria high pressure units and, finally, (4) non-metamorphic Piedmont-Liguria units that are only preserved along strike in the area

of the Chenaillet near Briançon (Manatschal et al. 2011). We attribute the Chenaillet ophiolites to that part of the Piedmont-Liguria Unit that occupied an upper plate position in the internal Western Alps (Schwartz et al. 2007) where, in our interpretation, the Austroalpine nappe system is totally lacking due to a westward lateral transition of the active margin of the Late Cretaceous to Paleogene Alpine subduction zone from the northern edge of the Adria continent into an intra-oceanic location (Molli 2008; Handy et al. 2010). The geometry of the shallow crustal parts of the Internal Alps depicted in this profile closely follows the geometries shown by the works of Henry et al. (1993), Michard et al. (2004), Lardeaux et al. (2006), and Ford et al. (2006), the surface-near parts of the Penninic Front exposing the Subbriançonnais and non-metamorphic Briançonnais units are drawn after a series of detailed profiles given by Gidon (1972).

The  $7.0 \text{ s}^{-1}$  velocity contour that runs parallel to and near the Moho depicted by controlled-source seismology in the Adria microplate rises SW-ward to a depth of some 15 km in the NE part of the profile beneath the HP Briançonnais. It is taken as marking the outline of the top of the Ivrea mantle wedge. The inferred overall geometry demonstrates that crust and mantle of the Adriatic microplate became wedged into the Alpine edifice and underplated shallow crustal layers in the internal parts of the Alps. The dip of the Periadriatic line is constrained to a depth of some 30 km by connecting the southern prolongation of the Periadriatic Line (Fig. 1), delimiting the eastern edge of the root of the Piedmont-Liguria units near the eastern margin of the Dora Maira massif at the surface, with the western front of the Ivrea mantle wedge paralleling the  $7.0 \text{ s}^{-1}$  contour. The exact geometry shown for the downward narrowing deepest parts of the internal Alps, i.e. the subduction channel separating the European and Adriatic (micro) plates, is basically unconstrained by geophysical data, and the details shown are merely conceptual. The downward bending of the  $7.0$  and  $7.5 \text{ s}^{-1}$  velocity contours across the suspected subduction channel is unable to depict the geometry of the subduction channel, due to the high velocities expected for the rocks within the subduction channel undergoing high-pressure metamorphism, velocities expected to be similar to those of the European lower crust, and at greater depth, similar to those of the adjacent mantle rocks.

The Oligocene age of formation of the steeply dipping root of the Piedmont-Liguria Ocean, associated with rock uplift associated with exhumation (Fox et al. 2016) of these high-pressure units along the front of the adjacent Adria microplate can be inferred from the cooling history of the Dora Maira high pressure unit evidencing rapid cooling from  $550 \text{ }^\circ\text{C}$  to some  $260 \text{ }^\circ\text{C}$  within an extremely short time interval between 32 and 30 Ma according to the pressure–time path constrained by petrological data in

combination with cooling ages derived from isotopic dating and fission track analysis (Gebauer et al. 1997; Rubatto and Hermann 2001; Berger and Bousquet 2008). The earlier stages of this exhumation are possibly related to exhumation within the subduction channel, triggered by the buoyancy of continental rocks at mantle depth (Chemenda et al. 1995). These time constraints indicate that both the formation of the steeply hinterland-dipping Penninic Front and the wedging of the Adria mantle and crust associated with backthrusting of the internal Alps over the Tertiary Piedmont Basin are contemporaneous. Both appear to be associated with the WNW-directed indentation of the Ivrea mantle during the earliest Oligocene (34–30 Ma).

The shallow NE crustal parts of the profile are covered by the Oligo-Miocene of the Tertiary Piedmont Basin and younger Plio-Pleistocene sediments. The construction of the profile is based on subsurface data acquired by AGIP (Pieri and Groppi 1981; Cassano et al. 1986; Schumacher and Laubscher 1996) that only vaguely constrain the geometry depicted. The  $5.5$ – $7.0$  velocity contours, as well as the Moho depth indicated by controlled-source seismology, indicate that within the Adria microplate Moho, lower crust and basement-sediment interface are E-dipping.

In the immediate vicinity of the Periadriatic line the thickness of the Tertiary Piedmont Basin along the profile is considerable. Carrapa and Garcia-Castellanos (2005) showed that the subsidence of the Tertiary Piedmont Basin is controlled by far-field compression and thrust loading rather than by extensional tectonics. Hence, the early stages of thrust loading are probably due to the Early Oligocene backthrust of the metamorphic units of the Western Alps onto the Adria microplate.

However, at the eastern end of the profile the Tertiary Piedmont Basin overlies “Apenninic” units (Fig. 2a). Compressional tectonics continued into the Miocene when the Ligurian Alps became involved in N-directed thrusting, together with Apenninic units exposed in the northern Monferrato hill (Fig. 1) west of Torino (Piana 2000), first during the so-called Paleo-Apenninic phase (Schumacher and Laubscher 1996) sealed by a Mid-Miocene unconformity. Later, Neo-Apenninic thrusting formed the morphologically visible thrust front north of the Torino and Monferrato hills. It is this N-directed Apenninic thrusting over the Adria microplate that also involved parts of the former Alps referred to as “Apenninic” units in the profile of Fig. 2a (see later discussion on the Alps–Apennines transition). Differential N-directed thrusting of joint Apenninic and Ligurian Alpine units calls for a sinistral strike slip zone that separates the joint Apennines-Ligurian Alps unit moving to the north from those parts of the Piedmont-Liguria units that are immediately adjacent to the Dora Maira backthrust, such as depicted near the eastern end of the section (Fig. 1a). The exact 3D geometry of this complicated Early Miocene to recent scenario still

remains largely unresolved and is known as the problem of the “Ligurian knot” (Laubscher et al. 1992).

## 2.2 Pelvoux profile (Fig. 2b)

The western part of this profile crosses the Vercors massif (Chaînes Subalpines) and the more internal Belledonne–Grandes Rousses–Oisans external massifs and their Mesozoic cover (Philippe et al. 1998; Dumont et al. 2008) (Fig. 1). Profile balancing by Bellahsen et al. (2012), taking into account the effects of normal faulting preceding Alpine orogeny (e.g. de Graciansky et al. 1989) related to the opening of Alpine Tethys, the total amount of Miocene shortening in these external parts of the Alps (Dauphinois) amounts to some 28 km.

In the external parts of the profile the P-wave velocity contours within the European lithosphere again exhibit a higher velocity gradient between 6.5 and 7.5 s<sup>-1</sup> near the Moho. However, the 6.0 s<sup>-1</sup> contour rises to an unusually shallow depth of <10 km twice, a feature that is considered robust and points to the presence of high-velocity material at shallow depth. The more external amongst the two anomalies is found underneath the Belledonne and Grandes Rousses massifs, an area that is known for the occurrence of the pre-Alpine Chamrousse ophiolitic complex (Guillot and Ménot 2009), associated with ophiolites and eclogites being part of a former Variscan suture zone. This suture zone evolved into a major SW-NE striking and ca. 1500 km long dextral wrenching zone in Late Carboniferous times, marking the SE margin of the European Variscides with Gondwana. This broad mega shear zone is known as the External Massifs Shear Zone (Guillot and Ménot 2009) or East Variscan Shear Zone (Padovano et al. 2012). The interpretation of this anomaly in terms of high-velocity material constituting this steeply inclined mega shear zone is admittedly speculative. The origin of the second more internal anomaly is unknown but possibly also related to velocity inhomogeneities related to pre-Alpine structures.

At the surface the Penninic Front is exposed as a spectacular thrust zone behind the Pelvoux massif and in the footwall of the Massif du Combeynot, dipping with ca. 35° to the E and associated with top-WNW out-of-sequence thrusting (Butler 1992; Ceriani et al. 2001; Ceriani and Schmid 2004; Trullenque et al. 2006). The linear extrapolation of this dip observed at the surface to depth is unconstrained by currently available geophysical data along this profile. Figure 2b opts for an extrapolation of the Penninic Front to great depth such as to join the subduction channel at the base of the Ivrea mantle wedge, consistent with what was observed along the Argentera profile described above. The 6.5 and 7.0 s<sup>-1</sup> contours that followed the lower crust of the European lithosphere in the

external Western Alps start to rise to shallower levels at an angle of around 45° after having crossed the Penninic Front, indicating that westward they enter high-pressure Briançonnais units.

The shallow crustal parts of the metamorphic core of the Alps are drawn according to data published in Barféty et al. (1995), Caby (1996), Ganne et al. (2006), and Lardeaux et al. (2006). In the Briançonnais units sedimentation continued until the Middle (internal Briançonnais) to Late Eocene (external Briançonnais) (Jaillard 1999) and hence predates thrusting at the Early Oligocene Penninic Front. They very much resemble those found along the Argentera profile. Again, almost the entire section across the inner Western Alps, except the very frontal part, is characterized by late-stage backfolding of pre-existing nappe contacts. The geometry of the top of the Ivrea mantle wedge is constrained by the 7.0 s<sup>-1</sup> velocity contour and controlled seismology data. Exact location and width of the subduction channel remain unknown, but an inverse velocity gradient of the 7.5 s<sup>-1</sup> contour associated with a down-bending of the same contour line points to the existence of a subduction channel below the Ivrea mantle wedge. A similar, although not identical geometry of a geological transect located near the Pelvoux profile was proposed by Zhao et al. (2015), who also postulated subduction of European continental lower crust all the way below the Ivrea mantle wedge, based on their interpretation of seismic imaging by P-wave receiver function analysis. In comparison, their model proposes a much shallower depth for the top of the Ivrea mantle wedge (at around 10 km) and a much wider (about 30 km) subduction channel below the Ivrea mantle wedge, features that are not compatible with the 3-D velocity model of Diehl et al. (2009) and the Moho depth as compiled by Spada et al. (2013).

The eastern parts of the profile are very similar to what was described for the Argentera section. They cross the northern parts of the Dora Maira massif and exhibit the same overturned root of the ophiolitic series of the Piedmont Lower Plate unit, juxtaposed with the front of the Adria microplate including the Ivrea mantle wedge along a backthrust onto the Tertiary Piedmont Basin, formed in the earliest Oligocene, i.e. contemporaneous with the formation of the Penninic Front. At its easternmost end the profile also reaches the Apenninic units thrust northward and perpendicular to the trace of the profile towards the Torino Hills in the Late Miocene.

## 2.3 ECORS-CROP profile (Fig. 2c)

This transect follows the trace of the geophysical campaign launched in 1985 by the French-Italian Étude Continentale et Océanique par Reflexion et Refraction Sismique-Progetto Strategico Crosta Profonda (ECORS-CROP)



geophysical campaign in the Western Alps involving high-resolution deep seismic sounding (Roure et al. 1996). The geological interpretation of this profile is a modified version of the profile previously published by Schmid and Kissling (2000).

The external Alps exhibit 27 km shortening, an amount very similar to that along the Pelvoux profile, calculated on the basis an area balance within an originally 17 km thick upper crust, assuming decoupling at the interface with the lower European crust. This shortening is of Miocene age and postdates thrusting at the Penninic Front (Fügenschuh and Schmid 2003). Note that Butler (1983) postulated very substantially higher amounts of shortening (>50 km) since his line balance assumes detachment along a floor thrust located only 1 km below base Triassic. The  $6.5 \text{ km s}^{-1}$  velocity contour beneath the external massifs, in the case of this profile the Belledonne Massif, indicates the presence of a steep tabular high-velocity body in the lower part of the upper crust that we again (see description Pelvoux profile) tentatively attribute to the East Variscan Shear Zone of Late Variscan age (Padovano et al. 2012). The velocities within the rest of the European upper crust are, apart from this anomaly,  $<6.5 \text{ km s}^{-1}$  all the way back to the very internal parts of the Alps below the Gran Paradiso massif, built up the high pressure Briançonnais unit. This new finding about the P-wave velocity in the upper crust of the external part of the Western Alps (Diehl et al. 2009) is incompatible with the presence of an allochthonous wedge of lower crust underneath the external Western Alps as postulated by Schmid and Kissling (2000) on the basis of gravity arguments (Bayer et al. 1989) and W-dipping seismic reflectors (Thouvenot et al. 1996) at the front of that putative wedge. Below the Gran Paradiso massif the  $6.5 \text{ km s}^{-1}$  contour steeply raises to a depth of only some 10 km, across the Penninic Front and into the area of the high-pressure part of the Briançonnais units, indicating a high-pressure metamorphic overprint also at great depth.

At the surface the profile crosses the type locality of the top-WNW Penninic Front (“Roselend Thrust” of Loprieno et al. 2011). This thrust, together with a 10 km wide belt of units attributed to the Valaisan paleogeographic domain, containing, amongst other lithologies, black schists inter-layered with mafic sills and low-T eclogites (Bousquet et al. 2002, 2012a; Loprieno et al. 2011) forms a planar band of strong reflectors that can be traced down to a depth of some 15 km along the ECORS-CROP reflection seismic profile (Thouvenot et al. 1996). Following Schmid and Kissling (2000), and in analogy with the Pelvoux profile discussed above, we kinematically link the Penninic Front with the base of the Adria mantle wedge.

The near-surface geometry of the upper crustal structures within the Briançonnais units, namely the non-metamorphic Zone Houllière and the High-Pressure

Briançonnais units (Ruitor, Zona Interna and Gran Paradiso nappes), as well as the geometries of the Piedmont-Liguria Units (Zermatt-Saas and Combin Zones) are drawn following Schmid and Kissling (2000) and Bucher et al. (2003, 2004). They are characterized by large-scale post-nappe backthrusts and backfolds formed between 35 and 31 Ma, i.e. contemporaneous with thrusting along the Penninic Front. High-pressure metamorphism related to subduction was dated at 48–47 Ma in the Briançonnais units, overprinted by greenschist facies conditions related to nappe stacking at around 43–39 Ma (Villa et al. 2014). Pressures associated with the high-pressure event increase eastward and reach eclogite facies conditions in parts of the Gran Paradiso internal massif, characterized by P-wave velocities around 6.0 to almost  $7.0 \text{ km s}^{-1}$  at greater depth according to the velocity model of Diehl et al. (2009).

Compared to the Briançonnais units high-pressure metamorphism within the more internal Zermatt-Saas-Fee unit (Piedmont-Liguria Lower Plate of Fig. 2c) and within small continental slices at its top indicates an earlier onset of a long and protracted high P overprint (62–42 Ma) according to age dating (e.g. Dal Piaz et al. 2001; Skora et al. 2015; Weber et al. 2015; Fassmer et al. 2016). The most internal parts of the ECORS-CROP profile cross the Sesia Zone, which together with the Dent Blanche Klippe, was classically considered as an “Austroalpine” unit. This is because the protoliths forming the Sesia Zone are undoubtedly derived from the Adria microplate in a paleogeographical sense (Compagnoni et al. 1977). However, in a plate tectonic context, the Sesia Zone that reached eclogite facies conditions with up to 2 GPa pressures in the period 85–60 Ma (Regis et al. 2014) and consisting of thin crustal sheets (Giuntoli and Engi 2016) is interpreted as having been accreted as a part of the former Lower Plate to the Upper Plate Units (i.e., the Canavese, Ivrea Zone and Southern Alps) during Late Cretaceous to early Cenozoic orogeny in the Western Alps, associated with the subduction of the Piedmont-Liguria ocean (Babist et al. 2006; Handy et al. 2010). The situation is different in the Austroalpine nappes proper where high-pressure metamorphism is significantly older and related to Cretaceous intra-continental subduction that evolved within the future Austroalpine units that formed the upper plate during Cenozoic orogeny (Schmid et al. 2004). In summary, the Sesia Zone is regarded as a lower plate unit following Froitzheim et al. (1996; their Figure 2) and Pleuger et al. (2007; their Figure 14) who proposed that the Sesia Zone represents a continental fragment floating within the Piedmont-Liguria oceanic domain that rifted and drifted away from Adria during the opening of Alpine Tethys in the Mesozoic. Two important geological observations support this finding. Firstly, the southern edge of the mostly continental Sesia high-pressure unit, immediately



adjacent to the Insubric Line, contains a high-pressure slice containing omphacite-bearing mantle peridotites, which are testimony of a former oceanic tract SE of the Sesia Zone that underwent high-pressure metamorphism (Pognante 1989; Barnes et al. 2014). Secondly, the non-metamorphic Canavese Zone, immediately south of the Insubric Line and north of the Ivrea Zone contains, amongst other lithologies, outcrops of exhumed mantle rocks overlain by Jurassic age radiolarites (Beltrando et al. 2015a), which points to the existence of an oceanic tract between the largely continental Sesia high-pressure unit and the Ivrea Zone. It is only the Canavese Zone, together with the Ivrea Zone and the Southern Alps that form the upper plate of the Alpine orogen in the internal parts of this transect. As demonstrated in the pioneering work by Elter et al. (1966) the Canavese Zone has to be considered the root of the highest nappe exposed in the Préalpes Romandes, the ophiolite-bearing nappe des Gets (see profile of Fig. 2d; unit labeled as Piedmont-Liguria Upper Plate). Both these units represent partly oceanic series that once formed the presently largely eroded upper plate of the Alpine edifice.

In contrast to the more southerly-located profiles, very thin slices of the Ivrea mantle wedge, accompanied by lower crustal lithologies of the Ivrea Zone, reach the surface of the earth, slices that are too thin to have a notable effect on the velocity contours. The main part of the Ivrea mantle wedge is confined to a depth  $>10$  km as indicated by the  $7.0 \text{ km s}^{-1}$  contour line. Along this profile the  $7.25 \text{ km s}^{-1}$  contour line reveals an inverted velocity gradient, allowing for localizing the base of the Ivrea mantle wedge above the subduction channel.

The Adria lower and upper crustal units are affected by backthrusts representing the easternmost extension of the Southern Alps. Details are drawn according to the interpretation given by Roure et al. (1990). This thrusting, detected in the subsurface of the Po plain in the course of the ECORS-CROP reflection seismic campaign, is of pre-Burdigalian (pre-20.5 Ma) age and affects part of the Gonfolite Lombarda sediments, providing an age range for thrusting of Late Oligocene to Early Miocene. Note that this profile does not reach the front of the N-directed Apenninic thrusts, which stays south of the SE end of this section.

#### 2.4 NFP-20 West profile (Fig. 2d)

This transect follows the trace of a Swiss geophysical campaign (NFP 20 “Deep Structure of Switzerland” whose results were published in Pfiffner et al. 1997). Earlier geological interpretation of the deep structure revealed by controlled source reflection and refraction seismology such as those by Escher et al. (1997) and Schmid et al. (2004) have been revised in the light of the 3D velocity model of Diehl et al. (2009).

Along this profile the effects of the foreland propagation of Alpine shortening induced the decollement of the entire Molasse basin along Triassic evaporite horizons, which led to the formation of the Jura Fold-and-Thrust Belt. The Alpine thrust front jumped by some 100 km to the north after about 12 Ma ago; this decollement is associated with some 26 km of shortening and roots in the external massifs (Aiguille Rouge and M. Blanc massifs) of the Alps (Burkhard and Sommaruga 1998). However, the total amount of shortening affecting the upper European crust, which along this transect also includes the lowermost Penninic nappes behind the Penninic Front, amounts to some 50 km according to an area balance along this profile, again assuming decoupling at the interface between upper and lower crust. This is almost twice as much as deduced for the ECORS-CROP profile but, in this case, also includes all the shortening that occurred since 35 Ma ago. The deep structure of the External Alps along this profile is characterized very marked undulations of the velocity contours between  $7.5$  and  $6.0 \text{ km s}^{-1}$ . Some of these undulations may again have been caused by velocity gradients related to pre-Alpine structures. However, one of these undulations, an antiformal structure located underneath the external massifs, appears to also affect the Moho; the closely spaced  $7.5$ ,  $7.0$  and  $6.5 \text{ km s}^{-1}$  velocity contours all consistently indicate an antiformal structure affecting the Moho and lower crust. A local up doming of the Moho by about 5 km correlates well with earlier observations in reflectivity pattern along the NFP20West near-vertical reflection profile (Pfiffner et al. 1997) that at the time were not further interpreted. As this Moho geometry has now been consistently observed by two independent seismic data sets, we assume this antiformal Moho-structure to be real and possibly representing a pre-Alpine feature.

For constructing the near-surface structures in the most external parts of the profile we mainly used Sommaruga (1997) for the Jura Mountains and Sommaruga et al. (2012) for the Molasse basin. In the case of the External Massifs, the Helvetic nappes and the Préalpes Romandes we based our interpretation of data provided by Burkhard (1988), Marchant (1993), Steck et al. (1997), Escher et al. (1997) and Burkhard and Sommaruga (1998).

The Penninic Front along this profile coincides with the south dipping Rhone-Simplon line, a wide dextral shear zone exhibiting initially ductile deformation starting to be active at around 35 Ma (Steck and Hunziker 1994; Schmid and Kissling 2000), which graded into brittle deformation from the Miocene onwards (Cardello and Mancktelow 2015). The Simplon normal fault located immediately east of this transect (Mancktelow 1992; Grasemann and Mancktelow 1993) represents a tensile bridge that accommodates orogen-parallel extension during the backthrusting over the Ivrea mantle wedge towards ESE and associated

top WNW thrusting at the Penninic Front of the Western Alps (Keller et al. 2006). The downward projection of the Rhone-Simplon line to some 15 km follows that proposed by Escher et al. (1997). At a >15 km depth any interpretation of the geometry of this fault and adjacent units is only loosely constrained by migrated reflection seismic data (Marchant 1993) and strike-parallel lateral projection of near surface structures. Our interpretation differs from that of Escher et al. (1997) at a depth >35 km because we conceive the Rhone-Simplon line as delimiting the northern border that acted as a lateral ramp for that part of the Alpine upper crust that moved towards WNW together with the Ivrea mantle wedge during its indentation.

The internal units of the Alps SE of the Rhone-Simplon line are characterized by spectacular backfolds within the Briançonnais units whose geometry down to a depth of about 20 km is constrained by reflection seismology (Marchant 1993; Steck et al. 1997) and orogen-parallel axial projections of surface structures (Escher et al. 1997; Keller and Schmid 2001; Kramer 2002). At a depth >20 km the P-wave velocity contours of Diehl et al. (2009) served as a guideline for constraining the geometry. The 6.0 km s<sup>-1</sup> contour indicates a large-scale velocity inversion that follows the SE limb of the Monte Rosa backfold down to a depth of some 30 km. There the 6.0 km s<sup>-1</sup> contour swings back into a region characterized by a normal velocity gradient and runs parallel to the 6.5 and 7.0 km s<sup>-1</sup> velocity contours that are typical for the European lower crust. As a consequence we limit the area characterized by such backfolding to the uppermost 35 km of the crust.

Farther to the SE the 6.0 and 6.5 km s<sup>-1</sup> contours rise to a shallow depth of only some 10 and 15 km, respectively. This rise occurs within the Sesia Unit that is known to exhibit eclogite facies conditions in its most internal parts only (Bousquet et al. 2012a). Because this eclogite facies part of the Sesia Zone is immediately adjacent to the Ivrea mantle rocks, the limit between the Adria mantle wedge and the under-plated Sesia zone is ill constrained. The near-surface structures of the Southern Alps are drawn after Cassano et al. (1986).

## 2.5 Ticino profile (Fig. 2e)

The Ticino profile runs parallel to the reflection seismic lines C1, C2 and C3 of NFP 20 project (Pfiffner and Heitzmann 1997); the southernmost parts of this profile are located near lines S4 and S5 of the same project (Schumacher 1997). At depth, the profile crosses the NE-most part of the Ivrea mantle wedge, very near its termination (see Fig. 3). Note also that this profile crosses a pronounced along-strike axial culmination (the core of the Lepontine dome; Berger et al. 2005; Berger and Mercogli

2006; Steck et al. 2013) at a location where the distance between external massifs and Periadriatic line is near a minimum (Fig. 1), going hand in hand with a maximum amount of shortening and exhumation. Since this transect is located east of the Rhone-Simplon line, dextral transpression allowing for top WNW thrusting along the Penninic Front is taken up by the Periadriatic line only.

According to an area balance, again assuming decoupling at the interface between upper and lower crust, the total amount of shortening affecting the upper European crust (including the Penninic nappes below the Valaisan and Briançonnais units) since around 35 Ma increases by as much as some 115 km when compared to the 50 km determined along the NFP20-West traverse. Rosenberg and Kissling (2013) measured some 71 km shortening along a profile located between our NFP20-West and Ticino profiles (their Simplon profile). This indicates a steady increase in the amount of post-collisional (post 35 Ma) shortening within the accreted European lower plate towards northeast into the Ticino transect.

In the external part of the profile the 6.0 km s<sup>-1</sup> contour rises to shallow depth underneath the frontal part of the external massifs, hinting at the existence of a pre-Alpine structure (External Massifs Shear Zone of Guillot and Ménot 2009) also along this profile. The near-surface features in the foreland of the Molasse Basin are drawn according to Sommaruga et al. (2012), the geometry of the external Aar and Gastern massifs is modified after Müller (1938) and Pfiffner (2011), the structures in the Helvetic nappes, overlying Penninic nappes and the Gotthard nappe (formerly considered a “massif”) are drawn after Menkveld (1995) and Pfiffner et al. (2010).

The internal parts of the transect, depicting the nappe pile within and above the Lepontine dome, are characterized by a large-scale backfold associated with the Northern Steep Belt located within the southern border of the Subpenninic Gotthard nappe (Schmid et al. 2004; Berger et al. 2005), separated by a flat-lying nappe pile from the Southern Steep Belt as defined by Milnes (1974), who was first in realizing the importance of post-nappe recumbent mega-folds affecting a previously existing nappe stack consisting of Briançonnais, Penninic and Subpenninic units. The formation of the Southern Steep Belt, the southern limb of a second more southerly located backfold, is closely associated with backthrusting of the Alpine nappe stack over the Ivrea mantle wedge and overlying crust in the area of the Southern Alps and along the Periadriatic Line that initiated between 35 and 30 Ma in a dextrally transpressive scenario (Schmid et al. 1987, 1989; Steck and Hunziker 1994). Note that Miocene thrusting in the Southern Alps is younger, i.e. post-20 Ma (Schönborn 1992). Based on the offset of the Alpine grade of metamorphism across the Periadriatic Line, the vertical

component of dextral transpression amounts to some 20 km along the trace of this profile. The amount of the simultaneously occurring horizontal displacement across the transpressive Periadriatic Line remains unknown due to the lack of markers; estimates vary between 100 km (Schmid and Kissling 2000) and 240 km (Handy et al. 2010). Details concerning the shallower parts of this central portion of the profile are drawn based on structural information by Milnes (1974), Huber et al. (1980), Leu (1987), Berger et al. (2005), Steck et al. (2013) and own observations. Reflection seismic data along line C1 and C2 (Pfiffner and Heitzmann 1997) were used for placing nappe contacts at the base of the Antigorio nappe and the Gotthard nappe, respectively, and for estimating the depth of the top of the European lower crust.

At a depth  $>20$  km the NW edge of the Ivrea mantle wedge is defined by the  $7.0 \text{ km s}^{-1}$  contour. At a depth of  $<20$  km the Periadriatic fault, whose dip is evaluated on the basis of surface observations and a series of reflection seismic profiles (Schmid and Kissling 2000; their Figure 5) delimits the NW edge of the Ivrea mantle wedge, including the Ivrea Zone that represents the adjacent Adriatic lower crust. The Ivrea Zone, including some ultramafic bodies at its northern rim, has been rotated along a near-horizontal axis into a vertical orientation due to Alpine deformation under low-grade metamorphic conditions (Handy 1987; Schmid et al. 1987; Zingg et al. 1990). The Ivrea Zone is well known for representing a map-view section across lower continental crust (Fountain 1976). It is likely that the topmost parts of the Ivrea Zone and mantle became highly tectonized under semi-brittle conditions, as is schematically shown in Fig. 2e, allowing for this rotation.


The  $6.5 \text{ km s}^{-1}$  contour rises to a shallow depth of around 10 km above the Ivrea mantle wedge also in this section, crossing from the European lower crust across the subduction channel over the presumed top of the Ivrea mantle wedge. The northern limit of the Ivrea mantle wedge and the adjacent Ivrea Zone are constrained by downward extrapolation of the Periadriatic Line to a depth of 20 km; at greater depth the Periadriatic line delimits the northern limit of the Ivrea mantle wedge.

The profile across the upper crust of the Southern Alps is partly based on the geological interpretation of reflection seismic data gathered during the NFP-20 and CROP campaigns (Schumacher 1997; Schumacher et al. 1997; lines C3-south, S4, S5, S6, S7 and CROP 88-1) and partly on the subsurface information provided by Cassano et al. (1986). Furthermore, we based our interpretation also on the work of Bertotti (1991) and Schönborn (1992). The total amount of shortening within the Southalpine retro-belt amounts to 74 km according to section balancing in the area crossed by this transect (Schumacher et al. 1997). A large, although not well constrained part (between 24 and

43 km along a section east of Lake Como according to Schönborn 1992) of this shortening pre-dates the Late Eocene. This leaves between 31 and 50 km of shortening that occurred in the Middle to Late Miocene. The lower bound of this shortening estimate is more realistic in view of the overall geometry of the transect depicted in Fig. 2e. The reason for the upward excursion of the  $6.5 \text{ km s}^{-1}$  contour inside the Adriatic upper crust remains unknown; possibly it is related to heterogeneities formed during Variscan tectonics.

## 2.6 Horizontal section (Fig. 3)

To discuss the overall shape of the Ivrea mantle wedge and lateral variations in its geometry the five profiles described above are complemented with a horizontal section across the 3-D P-wave tomographic model of Diehl et al. (2009) at a depth of 20 km (Fig. 3). This horizontal section shows (1) a curved area of positive P-wave anomalies  $>10\%$  nicely mimicking the well-known positive Bouguer



**Fig. 3** Horizontal section across 3-D P-wave tomographic model of Diehl et al. (2009) at a depth of 20 km. *Yellow triangles* indicate seismic stations used for tomographic inversion, *crosses* indicate inversion grid nodes. Note that the velocity structure is shown as percentage change relative to the 1-D initial reference model rather than in terms the P-wave velocities shown in Fig. 2. *Bold black contours* include well-resolved regions covering most of the area depicted. Compare Fig. 1 for major tectonic units outlined: *RSL* Rhone-Simplon Line, *PL* Periadriatic Line, *PF* Penninic Front, *ACL* Argentera-Cuneo Line. Comparison with Fig. 2 shows that the entire area with P-wave anomaly  $>10\%$ , as well as a part of the area with P-wave anomaly  $>5\%$  between north of Locarno and Cuneo is within the Ivrea mantle wedge

anomaly associated with the Ivrea mantle wedge (Kissling 1984; Serva 2005), (2) an extension of the Ivrea mantle wedge across the Periadriatic Fault visible north of Locarno (in reality the wedge underlies the N-dipping Periadriatic Fault, a backthrust in its hangingwall see Fig. 2e), and, (3) a possible dextral offset of the central axis of the Ivrea anomaly between profiles ECORS-CROP and NFP-20 West that leaves no trace at the surface of the earth when consulting the geological maps of the area.

Interestingly, the central axis of the high-velocity area is located east of the Periadriatic Fault in the north and gradually changes over to lying west of the Periadriatic line in the south. This indicates increasing amounts of underthrusting of the mantle wedge beneath the Internal Western Alps towards south, culminating in what is seen in the Argentera section (Fig. 2a): a Periadriatic Fault that acted as a huge backthrust and severe modification of the suture between Europe and Adria (lower plate Piedmont-Liguria ophiolites) in the form of a backfold with the high-pressure Briançonnais Units in its core (Fig. 2a).

### 3 Multi-stage evolution of the arc of the Western Alps

The following discussion on the evolution of the arc of the Western Alps integrates the findings described above with a review of existing knowledge on the multi-stage evolution of the Western Alps concerning kinematic evolution and time constraints derived from field data. In view of the considerable difficulties in exactly dating metamorphic events using isotopic ages, particularly in the case of HP metamorphism (see review by Berger and Bousquet 2008), we also heavily rely on biostratigraphical constraints. We also present tectonic reconstructions using Gplates reconstruction software (Boyden et al. 2011) that have already been applied to parts of the Western Mediterranean and eastern realm adjacent of the Alps–Apennines orogenic system (van Hinsbergen and Schmid 2012; van Hinsbergen et al. 2014a; Advokaat et al. 2014).

#### 3.1 Exhumation of Ivrea Zone and underlying lithospheric mantle in the Mesozoic

The Ivrea Zone is well known for exposing a map-view section across lower continental crust of the Southern Alps that formerly was part of the Adriatic microcontinent (e.g. Fountain 1976; Handy 1987; Handy and Zingg 1991). It includes a series of small peridotite lenses at its northern rim, due to later tilting in the context of Alpine orogeny. Restoration of the pre-Alpine setting via backrotation of Ivrea Zone and adjacent parts of the Southern Alps adjacent to the Periadriatic Line reveals that the Ivrea Zone was part of the


distal passive continental margin that formed in connection with Tethyan rifting culminating at around 180–170 Ma ago (e.g. Bertotti et al. 1993). The Ivrea Zone and underlying Ivrea mantle wedge were located near the western margin of the distal continental margin of the Adria microplate that had a strike (Weissert and Bemoulli 1985) almost parallel to the present-day NNE-SSW strike of the Ivrea mantle wedge (Fig. 3). Based on a combination of P–T estimates of pre-Alpine parageneses and radiometric dating it was inferred that crustal thickness was locally reduced to 10 km or less in the area of the Ivrea Zone and adjacent upper crustal units (Handy et al. 1999). Hence the present-day slope of the Moho, rising from some 40 km to about 20 km revealed by the Moho-map of Spada et al. (2013), and, rising to some 15 km in Fig. 2c, d, is interpreted as inherited from passive margin formation in Mesozoic times. The Canavese area, coinciding with the ocean-continent boundary, from which the Sesia extensional allochthon separated during rifting (Froitzheim et al. 1996; Beltrando et al. 2015a, b), is only some 50 km away from the Ivrea Zone.

The shallow depth of the top of the Ivrea mantle wedge inherited from Mesozoic rifting has important consequences regarding rheology and strength of the Ivrea mantle wedge during Alpine orogeny. The brittle to crystal plastic transition of peridotite is estimated to be at around 600° according to most workers (e.g. Druiventak et al. 2011). Temperatures in the depth range of the Ivrea mantle wedge (15–40 km) are between 400° and 650 °C in the central area of the Alpine orogen according to the steady state kinematic model of Bousquet et al. (1997). Hence, most of the Ivrea mantle wedge is expected to be controlled by frictional strength around 1000 MPa at 15 km depth in a compressional regime under dry conditions, reaching excessive values at greater depth (Byerlee 1978). At its front the Ivrea mantle wedge is surrounded by upper crustal nappes (Fig. 2) whose rheology is likely to be controlled by viscous creep of large quartz-bearing volumes of rock. Extrapolation of flow stress data to geological strain rates, combined with paleopiezometry using recrystallized grain size, predicts flow stresses between of 10 and 100 MPa within the same 400° and 650 °C temperature range (Stipp et al. 2002). This indicates that the strength of the Ivrea mantle wedge exceeds that of quartz-bearing units at its front by at least one order of magnitude. This suggests that indentation of the Ivrea mantle wedge into the Alpine nappe stack probably played an important role in the formation of the arc of the Western Alps, as will be further discussed below.

#### 3.2 Top-NNW thrusting during sinistral transpression (Stage 1, pre-35 Ma)

The orientation of stretching lineations in combination with associated sense of shear criteria have been applied to





**Fig. 4** Three generations of stretching lineations and associated senses of shear around the arc of the Western Alps. *Red arrow*: stage 1 top N transport; *blue arrows*: stage 2 top WNW transport; *green*: stage 3 top SW transport; *arrow pairs* reflect the variation of the measured lineations; see text for further details. Numbered areas of measurements and literature sources are as follows: area 1, Valaisan and Penninic Front (Loprieno et al. 2011); area 2, Zone Houillère of the Briançonnais Unit (Bucher et al. 2004); area 3, Subbriançonnais and Penninic Front (Ceriani et al. 2001; Ceriani and Schmid 2004); area 4, Penninic Front (top NW) and basal decollement of Cenozoic

flysch of the Dauphinois (top SW) (Bürgisser and Ford 1998; Trullenque 2005; Trullenque et al. 2006); area 5, eclogite facies lineations (top N) and greenschist facies lineations (top SW) in M. Viso ophiolites (Philippot 1988, 1990); area 6, Latest Eocene "schistes à blocs" at the base of the Embrunais-Ubaye nappes (Trullenque 2005); area 7, Penninic Front N of the Argentera massif (Trullenque 2005); area 7, thrust transport constraints from the Dauphinois of the Digne thrust system (Lickorish and Ford 1998; Ford et al. 2006)

deduce tectonic transport directions in highly sheared rocks from the Alps (e.g. Malavieille et al. 1984; Platt et al. 1989). Along the arc of the Western Alps the orientation of the lineations is dominantly transverse to the strike of the orogen. The inferred tectonic transport smoothly changes from top NW in the area of Monte Rosa to top WSW in the area of the Dora Maira area (Malavieille et al. 1984). This apparently simple arrangement, if taken to indicate sense of movement, leads to obvious difficulties in geometrical and plate kinematic terms (Platt et al. 1989). The major difficulties in interpreting kinematic data based on stretching lineation analysis is that (1) stretching lineations may indicate the orientations of finite strain (e.g. Ramsay 1980) rather than shearing during a particular stage of a complex orogenic evolution and (2) that orientations of older lineations may have been reoriented by younger deformations (e.g. Walcott and White, 1998; Keller and Schmid 2001; van Hinsbergen and Schmid 2012). Hence, only careful

local studies addressing the polyphase structural evolution of particular areas can reveal a more complex kinematic evolution. The results of such studies are compiled in Fig. 4.

A first generation of lineations indicating top north transport is best documented in areas of the Western Alps located behind, i.e. east, of the Penninic Front (areas 1, 2 and 3 in Fig. 4 and references therein). In these areas the orientations of this first generation of inferred transport directions is related to pre-35 Ma nappe stacking. According to detailed analyses of structural superposition this first generation of lineations pre-dates thrusting along the Penninic Front initiating at around 35 Ma and hence is of Eocene or even earlier age (e.g. Bucher et al. 2003; Ceriani and Schmid 2004; Loprieno et al. 2011). However the orientations of these first generation lineations scatter considerably between top-NW and top-NE due to later re-orientation. In the more internal parts of the internal



Western Alps the preservation potential of the orientation of such early stretching lineations is low. In most places they are not discernable any more because of intense shearing during subsequent deformation during stage 2 (see below). In the more internal units of the Aosta Valley, for example, they are seen to scatter too much for inferring transport directions of this earliest phase (Bucher et al. 2004). Only in some parts of the eclogitic lower plate Piedmont-Liguria units is a former top NNW eclogite facies lineation preserved (area 5 in Fig. 4; Philippot 1988, 1990). Interestingly, also parts of the External Western Alps (Pelvoux massif and its cover) are known to have been affected by stage 1 to N to NW-directed deformation of pre-Priabonian age (Dumont et al. 2011).

Large-scale structural considerations (Schmid and Kissling 2000; Handy et al. 2010) and plate kinematic constraints (Fig. 5) indicate that thrusting and plate convergence were top N to NNW before 35 Ma. This implies a scenario of sinistral transpression in the Western Alps during the early stages of orogeny when the upper plate tectonic units below which the Alpine nappe stack formed (Adria lithosphere; oceanic in case of the Western Alps, continental in case of the Eastern Alps) moved towards N to NNW, presumably past that part of the European plate presently located in SE France, as first proposed by Ricou and Siddans (1986). Pre-35 Ma plate convergence was associated with nappe stacking and early stages of exhumation of parts of nappes buried to high-P (e.g. Schmid et al. 1996, their Figure 8). A point located at the northern tip of Adria (Monti Lessini) moved some 260 km to the NNW in respect to stable Europe between 55 and 35 Ma go (1.3 cm/y) according to the reconstruction shown in Fig. 5d–f, which assumes that Adria was part of Africa at this time, and used the Europe-Africa plate circuit of Seton et al. (2012). This is more than the 195 km (0.98 cm/y) of shortening calculated by Schmid and Kissling (2000) for the same time interval but considerably less than the estimate of Handy et al. (2010) who inferred 465 km shortening for the interval 35–67 Ma (1.45 cm/y), both these estimates being solely based on interpretations of the geological structure and metamorphic history of the Alps. According to Handy et al. (2010) top-N thrusting started as early as some 84 Ma ago, consistent with the plate circuit estimates for the onset of convergence based on Seton et al. (2012), and first led to high-pressure metamorphism in the Sesia Zone (Manzotti et al. 2014), predating high-pressure metamorphism within the lower plate units derived from the Piedmont-Liguria Ocean that started at around 60 Ma ago (Berger and Bousquet 2008).

In summary, peak pressures within the Western Alps were reached in a scenario of sinistral transpression along an original approximately N-S striking segment of the western Alps (Fig. 5d, e) during this first stage. Most of the


**Fig. 5** Kinematic restoration of the Alps–Apennines orogenic system using Gplates reconstruction software (e.g. van Hinsbergen et al. 2014a) holding Europe fixed; for easier orientation all time slices also show the present day position of the external massifs and Tauern window in the Alps, as well as the present-day position of northern Adria. *Arrowheads* indicate displacement from a given time slice to the next one, they are anchored at the position of a chosen set of key locations at the chosen time slice. *Red lines* coast lines and outlines of external massifs and Tauern Window in the Alps; *blue lines* outlines of the Argentera, Pelvoux and Tambo-Suretta nappes; *green area* horizontal section across the Ivrea mantle slice taken from Fig. 3. The present day position of numbered locations shown is as follows: 1 Sardinia; 2 Corsica; 3 Dora Maira massif (Briançonnais); 4 Gran Paradiso massif (Briançonnais); 5 Tambo-Suretta nappes (Briançonnais); 6 Oberammergau (front Austroalpine nappes); 7 Dimaro (eastern end Tonale Line); 8 Mauls (western end Pustertal Line); 9 Slovenske Konjice (eastern end Gailtal line); 10 Cuneo (S end of Ivrea wedge); 11 Brione (N end of Ivrea wedge); 12 Pavia (Adria microplate); 13 Monti Lessini (Adria microplate); 14 Lignano (Adria microplate); 15 W of Ancona (frontal thrust of Umbria-Marche Apennines); 16 S of Parma (frontal thrust of external Ligurides, Apennines); 17 offshore Balears at 25 Ma ago (N front of oceanic Adria microplate = Piedmont-Liguria lower plate)

exhumation of the eclogites to moderate crustal levels very probably also took place during this first stage of the evolution of the arc of the Western Alps.

### 3.3 Top-WNW thrusting and WNW-directed indentation of the Ivrea mantle slice (stage 2, 35–25 Ma)

A second population of stretching lineations is associated with deformation that post-dates stage 1 top-NNW thrusting according to structural overprinting criteria. The top-WNW lineations formed during this stage can reliably be attributed directly to thrusting along the Penninic Front in the case of locations 1, 3, 4 and 7 in Fig. 4 (references cited therein). There the inferred thrusting directions vary between azimuth 260° and 345° with a mean at around azimuth 300°. The measurements at locations 2 and 5 were taken in more internal positions with respect to the Penninic Front while those at location 6 are from the footwall of the Penninic Front (lineations in the “schistes à bloc formation” immediately below the base of the Embrunais-Ubaye nappe stack; Trullenque 2005). Given the similar spread in orientations shearing during stage 2 top-WNW thrusting is likely also for locations 2, 5 and 6.

Several independent sets of data provide dating constraints. <sup>40</sup>Ar–<sup>39</sup>Ar dating of phengites that formed syn-kinematically with shearing of basement rocks of the Pelvoux massif in shear zones along the Penninic Front or in its immediate footwall vary between 35 and 30 Ma with one younger age at 27 Ma (Simon-Labric et al. 2009; Dumont et al. 2012). This dating confirms earlier evidence for the onset of top WNW thrusting at around 32 Ma based on fission track dating (Fügenshuh and Schmid 2003).



Since the Penninic Front is associated with dextral shearing along the Rhone-Simplon line (Fig. 1) and dextral shearing in the Insubric mylonites (part of PL of Fig. 1 around Locarno) the dating of such dextral shearing provides

further timing constraints. The “ductile Simplon shear zone” of Steck et al. (2015), predating late stages of normal faulting at the Simplon normal fault (Mancktelow 1992), was active between 34 and 25 Ma ago as inferred from the

ages of the mantle-derived andesite and tonalite (32–29 Ma) and synkinematic crustal aplite and pegmatite intrusions in the southern steep belt between Domodossola and Locarno (Steck et al. 2013). Dextral shearing in the mylonite belt of the Insubric line, a segment of the Periadriatic Line (PL in Fig. 1) initiated at around 32 Ma ago based on radiometric age data from the eastern margin of the Bergell pluton (32–30 Ma; von Blanckenburg 1992). The deep-seated Bergell intrusion including its subsequent fast exhumation is closely related to a combination of backthrusting and dextral strike-slip movements accommodated by the greenschist mylonites of the Insubric line (Schmid et al. 1989; Berger et al. 1996). The top-WNW senses of shear in the “schistes à bloc” formation (Kerckhove 1969) at location 6 of Fig. 4, affected by top-WNW shearing during a first phase of deformation (Trullenque 2005), provides a stratigraphic age constraint; this chaotic mélange overlies the Priabonian-age flysch de Champsaur (Bürgisser and Ford 1998), and hence, its age of formation is considered to be latest Priabonian to earliest Oligocene (around 34 Ma). A similar age constraint is provided for the sinistral senses of shear observed at the western termination of the Argentera-Cuneo Line at location 7 of Fig. 4 that is kinematically connected with the Penninic Front (Trullenque 2005); these mylonites are found in immediate tectonic contact with the grès d’Annot of Late Eocene to Early Oligocene age that terminate the evolution of the Eocene Northern Alpine foreland basin in the southern Subalpine chains (Ford and Lickorish 2004).

The end of stage 2 shearing is not equally well constrained but certainly pre-dates the activity of the left-lateral Giudicarie transpressive belt starting to offset the Periadriatic line at around 23–21 Ma ago (Scharf et al. 2013). An end of stage 2 at around 25 Ma ago is compatible with the onset of the Paleo-Apenninic phase (Schumacher and Laubscher 1996) at about this time. Stage 2 certainly pre-dates the onset of rapid counterclockwise rotation of the Corsica-Sardinia block at 21 Ma ago (Gattacceca et al. 2007) that we link to stage 3 oroclinal bending discussed in the next chapter.

At the large scale thrusting at the Penninic Front is interpreted to be kinematically connected to dextral strike slip along the Rhone Simplon and Periadriatic Lines on the one hand, and to sinistral strike slip along the Argentera-Cuneo Line on the other hand. The orogen-perpendicular lineations in the Western Alps compiled by Malavieille et al. (1984) are probably related to stage 2 top-WNW shearing and associated backfolding that is pervasive within entire volume of rocks between Penninic Front and the Periadriatic Line between Ivrea and Cuneo; this shearing formed new lineations and/or reoriented older ones (Fig. 1). It is proposed that this pervasive shearing, linked to WNW-directed indentation of the Ivrea mantle

wedge, first proposed by Laubscher (1971), also took place during stage 2 (35 and 25 Ma). The amount of top WNW displacement along the Penninic Front is not well constrained. Schmid and Kissling (2000) estimated WNW-directed displacement of the Ivrea Zone to amount to around 100 km, (Handy et al. 2010) proposed some 200 km. The reconstruction in Fig. 5c yields 93 and 150 km for the SW and NE ends of the Ivrea mantle wedge, respectively.

The reconstruction shown in Fig. 5c depicts displacement vectors that diverge between Western Alps and Eastern Alps and hence are bound to induce orogen-parallel extension in the Alps that is known to also have started at around 35 Ma ago (Pleuger et al. 2008; Beltrando et al. 2010). The backfolding and backthrusting, which is so typical for the Western Alps (see profiles of Fig. 2 and accompanying text) and which affected a pre-existing nappe pile, also occurred during stage 2 (e.g. Bucher et al. 2003). Hence, it is also associated with the indentation of the Ivrea mantle wedge that underthrusts much of the internal zones of the Alps (see Fig. 2a–c). The spectacular backthrust over the Cenozoic cover of the Tertiary Piedmont Basin shown at the eastern end the profiles shown in Fig. 2a, b also formed during stage 2.

### 3.4 Oroclinal bending and top SW thrusting in the southern part of the Western Alps (Stage 3, 25–0 Ma)

The external Alps south of the Pelvoux massif are well known for late-stage top-SW thrusting (e.g. Fry 1989; Lickorish and Ford 1998; Bürgisser and Ford 1998) that is not seen farther to the north along the Western Alpine arc. By studying overprinting relationships south of the Pelvoux massif (area at location 4 in Fig. 4) Trullenque (2005) discovered that this top-SW thrusting actually post-dates stage 2 top WNW thrusting. Lickorish and Ford (1998) have shown that 10.5 km out of a total of around 21 km shortening, associated with minor thrusting rooting in the basement of the Argentera Massif, occurred in Late Miocene to Pliocene times. Ritz (1992), based on a paleostress analysis in the Digne, Castellane and Nice thrust systems, found that during Middle Miocene to recent times the  $\sigma_1$  trajectories varied from NE-SW in the Digne area, gradually turning into a N-S orientation in the Castellane arc and eventually to a NW-SE direction in the case of the Nice thrust sheet. This same radial arrangement was found for the p-axes derived from stress inversion of earthquake focal mechanism data by Delacou et al. (2004, their Figure 5). This radial pattern is very likely related to rotation around a pivot located east of the Argentera massif in the area of the “Ligurian knot” (Laubscher et al. 1992), i.e. in an area where the 3 Moho’s beneath Alps, Apennines and northern Tyrrhenian Sea meet (Spada et al. 2013).

Three largely independent lines of evidence support the idea that oroclinal bending in the southernmost Western Alps is to be seen in connection with the structural observations mentioned above. Firstly, Collombet et al. (2002) reported counterclockwise rotations about a sub-vertical axis in respect to stable Europe that increase from  $68^\circ$  in the Ubaye region (area between profiles 1 and 2 of Fig. 1) to  $117^\circ$  in the Ligurian Alps SE of Cuneo (Fig. 1). The data were collected in sedimentary rocks of the Briançonnais tectonic unit. The age constraints for magnetization are rather loose; the authors propose that the rocks were remagnetized after the backthrusting phase (our stage 2) and therefore in the Late Oligocene. Secondly, all the major tectonic elements of the internal Alps derived from the Briançonnais and Piedmont-Liguria paleogeographical domains can be followed into western Liguria and all the way to Genova and slightly beyond (Fig. 1). In the Ligurian Alps all the pre-Oligocene Alpine structures, sealed by the Tertiary Piedmont Basin, are top-S in present-day coordinates. Since the counterclockwise rotation of stable Adria only amounts to  $10 \pm 10^\circ$  (van Hinsbergen et al. 2014b) or  $20^\circ$  (Marton et al. 2011) at least a part of the  $49^\circ$  differential rotation of Liguria in respect to the Ubaye region ( $117^\circ - 68^\circ = 49^\circ$ ), namely  $39 \pm 10^\circ$ , is probably related to oroclinal bending during stage 3. It is likely that oroclinal bending during stage 3 was partially driven by eastward trench retreat of the Apennines slab. However, the rotation in the Ubaye region itself, located north of the Argentera-Cuneo Line is probably also due to stage 2 indentation of Adria. Thirdly, a  $50^\circ$  counterclockwise rotation phase of the Corsica-Sardinia block occurred between 20.5 and 16 Ma ago (Gattacceca et al. 2007) around a pivot near the N tip of Corsica, as Corsica and Sardinia drifted away from their pre-Miocene location south of southern France (Advokaat et al. 2014); this virtually demands substantial counterclockwise rotation in Liguria associated with oroclinal bending. Note that such counterclockwise rotation is directly demonstrated by the  $\sim 45\text{--}50^\circ$  anticlockwise rotation of mostly Oligocene sediments of the Tertiary Piedmont Basin sealing pre-Oligocene Alpine structures evidenced by paleomagnetic data (Maffione et al. 2008).

Oroclinal bending in the southernmost Western Alps during stage 3 is evidently associated with orogeny in the Apennines. Slab rollback of the Adria plate towards N in the northernmost Apennines is the underlying driving force for both oroclinal bending and the rotation of Corsica and Sardinia. The discussion of the Alps–Apennines transition, presented in the next chapter, will shed more light to the Miocene history of the Ligurian Alps that cannot be understood without discussing orogeny in the Apennines and its spatial and temporal relation to orogeny in the Alps.

## 4 The Alps–Apennines transition in space and time


### 4.1 Tertiary Piedmont Basin and Epiligurian basins

Late Eocene to Neogene sedimentary basins found in the interior of the Western Alpine arc, the Tertiary Piedmont Basin, and similar basins found on top of the External Ligurides of the Northern Apennines, referred to as “Epiligurian” basins, play a crucial role for understanding the spatial and temporal transitions between the west- and north-verging Alps and the east- and north-verging Apennines (Fig. 6).

The southern limit of the outcropping pre-Pliocene parts of the Tertiary Piedmont Basin (Fig. 1) transgresses the high-pressure lower plate units of the Ligurian Alps (i.e. the high pressure Briançonnais and Piedmont-Liguria units), as well as adjacent upper plate units of the Alps derived from the Piedmont-Liguria Ocean found immediately east of Genova (i.e. the Antola nappe). As first pointed by Elter and Pertusati (1973) the Antola nappe, predominantly consisting of Helminthoid Flysch, is best considered as a former part of the Alps; they wrote: “La falda del Monte Antola e le altre formazioni liguri comprese nel triangolo Genova-Val Staffora-Levanto sono da considerarsi appartenenti alle Alpi in quanta sono saldate a queste dai terreni trasgressivi del Bacino terziario Piemontese fin dall’Eocene”). While earlier works suggested that this basin initiated by subsidence in a scenario of crustal stretching (Mutti et al. 1995) there seems to be some consensus in more recent publications that the Oligocene to Lower Miocene (Aquitania) sediments of this basin were laid down in a compressive to transpressive setting (e.g. Carrapa and Garcia-Castellanos 2005; Mosca et al. 2010).

As shown in the profiles of Fig. 2a, b the westernmost parts of the Tertiary Piedmont Basin were installed onto the upper crust of the Adria plate, which in turn overlies the Ivrea mantle wedge. It formed the foredeep of the backthrust and -folded Western Alps during the Oligocene, which later became buried under post-Messinian cover. Such syn-sedimentary backthrusting to the east to southeast is seen in the seismic sections of lines 3 and 4 of Mosca et al. (2010, their Figure 4). Thrusting is sealed by the Burdigalian unconformity that is ubiquitous in seismic sections of the subsurface of the Po plain (e.g. Schumacher and Laubscher 1996 and references therein). Such pre-Burdigalian thrusting is also seen along the front of the Southern Alps in western Lombardy where it affects sediments of Oligocene to Lower Miocene age, the Gonfolite Lombarda Group *sensu lato* (e.g. Bernoulli et al. 1989, 1993) that are part of the Southalpine foredeep





**Fig. 6** Interpretative map sketch of the Alps–Apennines transition; see text for further discussion

(Garzanti and Malusa 2008) and laterally connect with the Tertiary Piedmont Basin (Mosca et al. 2010). Subsidence in the Piemonte region continued in Burdigalian and younger times, when this basin started to become involved in N-directed thrusting in connection with the Apenninic orogeny. The Torino and northern Monferrato hills immediately east of Torino (Fig. 1) are also considered as a part of the Tertiary Piedmont Basin by most authors although they would better be considered a part of the Epiligurian basins (see below). They morphologically mark the northern front of the Apenninic orogen that formed in Late Miocene to Pliocene times. These two northernmost parts of the Tertiary Piedmont Basin transgressed an "Alpine-type" basement in case of the Torino hills (Piana 2000) and a part of the External Ligurides in the case of the northern Monferrato hills (Laubscher 1971; Mosca et al. 2010; Festa and Codegone 2013). Together with the rest of the Tertiary Piedmont Basin that transgressed Alpine structures they became thrust northwards as wedge-top

basins, much like the so called Epiligurian basins of the External Ligurides, described below, along-strike farther to the east and southeast.

The term Epiligurian basins is used for wedge-top basins overlying the External Ligurides of the northern Apennines (Fig. 6) whose age-range and sedimentary characteristics are similar to those of the Tertiary Piedmont Basin (Molli et al. 2010 and references therein). The sedimentary sequence starts with the deposition of the deep marine Middle to Upper Eocene Monte Piano marls above a regional unconformity that indicates some pre-Apenninic deformation of parts of the External Ligurides that represent a fossil ocean-continent transition (Marroni et al. 2001) within the Adria plate that was part of the upper plate of the Alpine orogen. The Early Oligocene Ranzano formation was laid down above a second unconformity and received detritus from the metamorphic units of the Central Alps (Leontine dome) that started to become exhumed north of the Periadriatic line (Garzanti and Malusa 2008).



In the Early Oligocene the External Ligurides were part of the Adriatic foredeep in respect to the Southern Alps that formed due to “retroshearing” (Beaumont et al. 1994) within the Alpine orogen (Schmid et al. 1996). Subsidence and siliciclastic sedimentation in the Epiligurian basins continued during the Late Oligocene when these basins and the underlying units became involved in the Apenninic orogeny during its earliest stages. From ca. 23 Ma onward the Epiligurian basins evolved into wedge-top basins as the External Ligurides became underplated by the accretion of Subligurian and Tuscanide units within the evolving Apenninic orogen (Molli et al. 2010 and references therein). Very recently Piazza et al. (2016) provided evidence for intense syn-depositional sinistral transpression during Oligocene to Early Miocene times sealed by a Burdigalian unconformity within one of the Epiligurian basins. This sinistral transpression is contemporaneous with sinistral strike slip along the E-W striking Villalvernia-Varzi line (Elter and Pertusati 1973) that defines the boundary between the Antola nappe often considered as a part of the Alps in the south and the External Ligurides considered as a part of the Apennines in the north (stippled line in Fig. 1 north of Genova). This sinistral strike slip movement was active in Late Oligocene to Early Miocene times as well (Di Giulio and Galbiati 1998; Schumacher and Laubscher 1996). Laubscher (1991) considered the Villalvernia-Varzi line as accommodating the WNW-directed indentation of the Ivrea mantle wedge previously discussed (stage 2, during the evolution of the arc of the Western Alps). Although we consider the direct link between the sinistral Argentera-Cuneo line and the Villalvernia-Varzi line proposed by Laubscher (1991) as unlikely (see Figs. 1, 6) sinistral strike slip motion along the Villalvernia-Varzi line and sinistral transpression in the Epiligurian basins were probably kinematically connected with stage 2 WNW-directed indentation of the Ivrea mantle wedge during the Oligocene.

In summary, the Tertiary Piedmont Basin seals very substantial pre-Late Eocene Alpine structures. The Epiligurian basins are of a similar type and age but they were deposited above an angular unconformity onto a unit that never underwent substantial Alpine deformation, namely the External Ligurides of the Northern Apennines (Fig. 6). Later, i.e. from about 23 Ma onwards, the latter represented wedge-top basins riding on the External Ligurides, which thrust over more external units of the Northern Apennines towards NE (Molli 2008; Molli et al. 2010, and references therein). In essence, the stratigraphic base of these basins post-dates the pre-Late Eocene top-NNW thrusting during stage 1 sinistral transpression in the Western Alps described earlier while it pre-dates the onset of top-NE thrusting during the Apenninic orogeny, which is contemporaneous with stage 3 oroclinal bending and top

SW thrusting in the southern part of the Western Alps. It follows from this that substantial parts of these basins probably represented syn-orogenic deposits during stage 2 top-WNW thrusting and indentation of the Ivrea mantle wedge into the Western Alps at 35–25 Ma ago.

#### 4.2 The Alps–Apennines transition in map view

An interpretative map sketch of the Alps–Apennines transition is presented in Fig. 6. Considerable parts of this area are either buried by deposits of the Tertiary Piedmont Basin and very young Cenozoic cover of the Po plain or by the waters of the Ligurian and North Tyrrhenian Sea; hence this compilation in Fig. 6 remains speculative in many respects.

The continuation of the undeformed or little deformed Alpine foreland has to be located in Variscan Corsica that became thrust by Alpine allochthonous units until the Late Eocene (Malavieille et al. 1998; Molli 2008). Note, however, that the present-day NW–SE strike of the Late Eocene Alpine front across the Ligurian Sea and Corsica restores to a N–S strike after the retro-deformation of 47° post-Eocene rotation of the Ligurian Alps (Maffione et al. 2008; see Fig. 5d, e) and 50° rotation of Corsica-Sardinia (Gattacceca et al. 2007). The Alpine allochthons in Corsica consist of, from bottom to top, the high-pressure units of the Tenda “massif” representing formerly subducted and subsequently exhumed parts of Variscan Corsica that remained devoid of any Alpine overprinted in the west, the high-pressure units of the Piedmont-Liguria Ocean and finally overlying non-metamorphic ophiolitic and continental nappes (Balagne and Nebbio nappes). The two lower units clearly represent lower plate units of the Alps that have to be correlated with the Briançonnais and Piedmont-Liguria units of the Western Alps, respectively. Although the origin of the uppermost non-metamorphic units is controversial (see Marroni and Pandolfi 2003 and references therein) we follow Malavieille et al. (1998) who argue for an origin of these highest units from the ocean-continent transition at the eastern margin of the Piedmont-Liguria Ocean. This makes them comparable to the non-metamorphic upper plate units of the Western Alps mapped in Figs. 1 and 6 (Préalpes Romandes, Embrunais-Ubaye and Western Liguria Helminthoid flysch). Oligocene–Miocene extension and transtension localized in the Gulf of Lion and the Ligurian Sea strongly affected the entire Corsican belt (Fig. 6), reducing its crustal thickness to normal thickness. In map view the width of the Alpine belt in Corsica is drastically reduced in comparison with the Western Alpine orogenic belt. We interpret this reduction in width to be partly related to the switch in subduction polarity between Alps and Apennines that led to a new west-dipping “Apenninic” subduction zone that must have

evolved offshore eastern Corsica since latest Eocene or earliest Oligocene times (Fig. 6; Doglioni et al. 1998; Carminati et al. 2004; Molli and Malavieille 2011).

This new plate boundary was located east of the Corsica basin (Mauffret et al. 1999; see also CROP-03 and M-12A seismic sections in Finetti et al. 2001 and Finetti 2005). It evolved after cessation of Alpine, southeastward subduction in the Oligocene, and was followed by the two-stage opening of the Gulf of Lion back-arc basin. In a first, Oligocene stage, at around 30 Ma Corsica-Sardinia rifted off the Provence margin of southern France (Séranne 1999). This was followed by the 21–16 Ma Corsica-Sardinia rotation around a pole located somewhere between the northern tip of Corsica and the Ligurian coast (Advokaat et al. 2014). The new plate boundary probably had a ~N-S-direction, slightly oblique to the NNW-SSE present-day strike of the Alpine orogen in Corsica (Fig. 6). In respect to the Apenninic orogeny both lower and upper plate units of Alpine Corsica occupied an upper plate position, remaining largely unaffected by Apenninic shortening, while easterly and northerly adjacent parts of the former Alps were heavily reworked when backthrustured onto the Adria continental margin and subsequently stretched as the Adria plate below the North Apennines rapidly rolled back eastward (Grandjacquet and Haccard 1977; Doglioni et al. 1998; Rosenbaum and Lister 2004; Spakman and Wortel 2004; Marroni et al. 2010).

The northern margin of the lower plate metamorphic core of the Ligurian Alps (AFSZ in Fig. 1) is not exposed being buried beneath the Cenozoic cover of the Tertiary Piedmont Basin and younger cover (Fig. 1). In Fig. 6 it is mapped according to subsurface seismic data (Mosca et al. 2010). Along a transect located west of Genova (profile A in Mosca et al. 2010) the metamorphic units of the Ligurian Alps were thrustured northward by about 50 km over the Adria continental autochthonous during the Apenninic orogeny in Miocene-Pliocene times. This implies that the western continuation of the older (pre-25 Ma) Argentera-Cuneo Line, delimiting the southern boundary of the indenting Ivrea mantle wedge and overlying internal Western nappe stack during the stage 2 top-WNW thrusting in the arc of the Western Alps, must be cut off by a N-S striking sinistral shear zone (Figs. 1, 6). This putative shear zone, which is only schematically drawn in Fig. 6, delimits the western edge of the Ligurian Alps affected by these 50 km of Apenninic top N thrusting from a narrow strip of the Southern Alps retro-belt in the west.

Eastward, i.e. north of Genova, the northern limit of the metamorphic core of the Ligurian Alps bends into a N-S strike and is exposed at the surface forming the well-known Sestri-Voltaggio line (or zone; Cortesogno and Haccard 1984) that is often taken as representing the Alps–

Apennines junction (e.g. Castellarin 2001). Although reworked in later Neogene times (Crispini et al. 2009) this line basically represented a Late Paleocene to Early Eocene normal fault (Hoogerduijn Strating 1994) sealed by the deposits of the Tertiary Piedmont Basin. This normal fault juxtaposes the high-pressure rocks of the Voltri Group in the west (1.5–2.5 GPa), attributed to the Alpine lower plate, with low metamorphic grade units to the east (Sestri-Voltaggio Zone, various Ligurian flysch units and finally the Antola nappe; see Figure 8d in Molli et al. 2010), whose grade of metamorphism progressively decreases eastward and up-section (see Figure 2 in Capponi et al. 2009). It finally reaches the diagenetic zone in terms of illite crystallinity (Ellero et al. 2001) over a short horizontal distance of only some 5–7 km. Following Elter and Pertusati (1973) who considered the Antola nappe as an integral part of the former Alps that became backthrustured during Alpine orogeny, we attribute the units east of the Sestri-Voltaggio normal fault to the Alpine upper plate (Fig. 6).

The unknown Alpine-type basement of the Torino hills buried under the deposits of the Tertiary Piedmont Basin and the Antola nappe were mapped together as parts of a strip of former Alpine upper plate units mostly buried under Cenozoic deposits in Fig. 6. During the Apenninic orogeny, together with the metamorphic core of the Ligurian Alps, they thrustured the External Liguride units of the Monferrato hills along the Rio Freddo Fault Zone (Piana 2000) striking NW-SE between the Torino and Monferrato hills. The Rio Freddo Fault Zone links up with the E-W-striking Villalvernia-Varzi line (Fig. 1), a Late Oligocene to Early Miocene thrust, active during the deposition of a part of the Tertiary Piedmont basin fill, with a strong sinistral strike slip component (Laubscher et al. 1992; Schumacher and Laubscher 1996). This sinistral transpressional fault is responsible for laterally transporting the External Ligurides exposed in the Monferrato hills in front of the Antola nappe and the Torino hills Alpine basement. At its eastern termination the Villalvernia-Varzi line bends into a N-S-strike as it transforms into an Apenninic top NNE thrust of the Antola nappe over the external Ligurides in the north and over the Internal Ligurides further south (Fig. 6). The polyphase history of the Antola nappe, with early and Alpine top-NW thrusting and folding, followed by top-NNE Apenninic thrusting over the Ligurides, is well documented (e.g. Marroni et al. 1999, 2002a; Levi et al. 2006). Offshore the continuation of this thrust is only tentatively drawn in Fig. 6; southward it is interpreted to eventually abut the northern continuation of the west-dipping upper/lower plate boundary of the Apennines orogen.

The next eastward unit of the Apennines, the Internal Ligurides, have also to be considered as having represented an integral part of the former Alps. They occupied an upper

plate position before they became backthrust to the NNE during the Apenninic orogeny. The internal Ligurides are characterized by the presence of classical ophiolites and an Upper Jurassic to Lower Cretaceous sedimentary cover (cherts, Calpionella limestone and Palombini shales) associated with Upper Cretaceous–Paleocene turbiditic sequences (Marroni and Pandolfi 1996) that represent the pelagic cover of the ophiolitic basement of the Piedmont-Liguria ocean (Decandia and Elter 1972) whose stratigraphy is virtually indistinguishable from that of the ophiolites of the Alps *sensu stricto* (Bernoulli et al. 2003). Moreover, since the pioneering work of Grandjacquet and Haccard (1977) several authors evidenced classical structural overprinting patterns at all scales, clearly showing that the early tectonic history of the Internal Ligurides was characterized by a top W to NW Alpine tectonic fingerprint, associated with low-grade metamorphism reaching some 300 °C (Pertusati and Horrenberger 1975; van Wamel 1987; Marroni and Pandolfi 1996; Ellero et al. 2001).

The External Ligurides are far-travelled and detached units that are confined to the northernmost Apennines between Torino and Bologna (Fig. 6). They overlie thin slices of Subligurian units and the Cervarola unit representing flysch deposits of the Adriatic foredeep outcropping in two small windows (Molli et al. 2010). The External Ligurides dominantly consist of detached Cretaceous–Paleocene flysch sequences (Helminthoid flysch) overlying basal mélange-type complexes (Marroni et al. 1998, 2001, 2002b). Two main subgroups of units can be recognized: those associated with ophiolites and with ophiolite derived debris (Molli 1996), and others without ophiolites and associated with fragments of Mesozoic sedimentary sequences and conglomerates with continental Adria affinity often remarkably similar to those found in the Alpine upper plate units of the Préalpes Romandes (Elter et al. 1966). Hence, the External Ligurides are regarded as representing the former ocean-continent transition within the Adria plate. Very commonly the stratigraphic base of the Oligocene part of the Epiligurian basins only exhibits a modest angular unconformity up to few tens of degrees (Piazza et al. 2016). Due to the syn-depositional sinistral transpression during Oligocene to Early Miocene times sealed by a Burdigalian unconformity mentioned earlier, the base of the Burdigalian and younger deposits of the Epiligurian succession locally overlies overturned older sequences of Late Cretaceous to Early Miocene successions (Piazza et al. 2016). In summary, the External Ligurides represent an ocean-continent transition area within the Adria upper plate of the Alps that was only locally overprinted by substantial late Alpine deformations of Oligocene to Early Miocene age, largely synchronous with stage 2 deformation in the Western Alps.

A substantial Alpine overprint has not been documented in the southerly adjacent Tuscan Ligurides,

representing a Piedmont-Liguria ophiolitic sequence identical to that observed in the Internal Ligurides or the Alps (Principi et al. 2004). Due to massive extension locally leading to core complex formation (Brunet et al. 2000), Tuscany exposes deep portions of the Apenninic orogen that locally reached blueschist facies conditions with pressures up to some 1.3 GPa (Rossetti et al. 2002), mostly affecting the underlying Tuscan nappe that represents the lower plate unit during Apenninic orogeny. Locally, also the immediately overlying Tuscan Ligurides (Jolivet et al. 1998; Brunet et al. 2000; Rossetti et al. 2002) became involved in subduction during the Apenninic orogeny. Hence, it is likely that the Tuscan Ligurides only became involved in W-directed Apenninic subduction and suffered penetrative deformation within a W-dipping subduction channel. A significant ophiolitic complex that is part of the Tuscan Ligurides is exposed on Gorgona Island (Orti et al. 2002). Metasediments associated with these ophiolites suffered blueschist facies metamorphism reaching about 1.5 GPa at around 25 Ma (Rossetti et al. 2001). Gorgona island is located far offshore Tuscany (Fig. 6) and very close to the northern continuation of the new plate boundary that formed during the earliest stages of Apenninic orogeny associated with W-directed subduction and subsequent roll back of the mantle slab below the North Apennines (Spakman and Wortel 2004). W-directed subduction initiated east of the Corsica basin that represents a >8.5 km thick rift basin filled with undeformed Late Oligocene to Burdigalian sediments (Mauffret et al. 1999; Finetti 2005) deposited as the Adriatic slab started to roll back, causing extension in the Corsica basin located in the Apenninic upper plate. The range of radiometric ages obtained for the high-pressure metamorphism in the Tuscan units is also significantly younger (26–20 Ma; Rossetti et al. 2002) when compared with the radiometric ages obtained in the ophiolitic units of Corsica where Cretaceous to latest Eocene ages (Vitale Brovarone and Herwartz 2012, and references therein) have been obtained. In summary, the Tuscan Ligurides are very likely to only have been affected by Apenninic orogeny, associated with W-directed subduction in latest Oligocene to Miocene times.

The Mesozoic Adriatic passive margin stratigraphy that accreted to the overriding European plate (Corsica) and the Ligurides during the Apenninic orogeny is exposed in the Tuscan nappe. The Chattian–Aquitainian Macigno turbidites (Di Giulio 1999) represent the youngest sediments involved. This 2.5–3 km thick turbiditic sequence is testimony of the deposition of siliciclastics derived from the Alps, particularly from the Lepontine dome (Garzanti and Malusa 2008). This dome shed an immense volume of metamorphic basement estimated at  $17\text{--}20 \times 10^3 \text{ km}^3$  (Di

Giulio 1999) as the Lepontine dome was uplifted by retro-thrusting and exhumed by erosion. This led to deposition of the Macigno pile of turbidites onto the still undeformed Adriatic foredeep during retro-folding and thrusting of the Alps north of the Periadriatic Line, predating most of the backthrusting of the Southern Alps in Lombardy (Schmid et al. 1996). Related exhumation, erosion and re-deposition onto the still undeformed Adriatic continental margin are largely synchronous with stage 2 top-WNW thrusting and WNW-directed indentation of the Ivrea mantle wedge (Stage 2, 35–25 Ma).

The more external Apennines thrust sheets involve flysch deposits of the foredeep only. The Cervarola thrust sheet is made up of Aquitanian to Early Langhian turbidites that still received detritus from the Alps. The more external Umbria-Marche thrust sheets involve the Marnoso Arenacea sequence of Serravallian to Tortonian age (Di Giulio 1999). The post-Messinian frontal imbricates are buried below younger deposits of the Po plain (Picotti and Pazzaglia 2008).

## 5 Lateral or temporal change in subduction polarity between Alps and Apennines? A discussion

Since the early days of Argand (1924) most authors (e.g. Mattauer et al. 1981; Durand-Delga 1984; Molli et al. 2006; Molli and Malavieille 2011; Handy et al. 2010) regarded the pre-Oligocene Corsican nappe stack as a continuation of the Alpine orogen. Some authors (e.g. Principi and Treves 1984; Lahondère 1996; Jolivet et al. 1998; Brunet et al. 2000), however, regarded Alpine Corsica as an integral part of the Apennines, both having formed in connection with a W-dipping subduction zone since Cretaceous times. They considered Variscan Corsica as a backstop of the Apennines orogenic wedge onto which the deepest part of the Apennines accretionary complex, namely Alpine Corsica, was backthrust. This view essentially transfers the area of the Alps–Apennines transition more to the north into the area around Genova, which is rather unlikely in the light of the available data on Alps and northern Apennines summarized in the previous chapters.

Debate also exists on whether the Alpine subduction zone with Eurasia in a lower plate position has continued beyond Corsica (e.g., Handy et al. 2010), or whether a lateral change in pre-Oligocene subduction polarity occurred in an area located somewhere between Corsica and Sardinia active since Late Cretaceous to Early Eocene times (e.g. Marchant and Stampfli 1997; Faccenna et al. 2003; Lacombe and Jolivet 2005; Lustrino et al. 2009; Argnani 2012; Advokaat et al. 2014; van Hinsbergen et al. 2014a). This transition is supposed to have occurred along a discrete trench–trench transform located SE of Corsica–

Sardinia (e.g. Argnani 2012) or more diffusively, with partly overlapping trenches associated with opposite subduction polarity somewhere offshore east of Corsica (e.g. Lacombe and Jolivet 2005). An often quoted argument in favor of this view is the existence of Late Eocene magmatism in Sardinia (Lustrino et al. 2009) within but a tiny fraction (Calabona subvolcanics) of the dominantly latest Oligocene and younger igneous magmatic province of Sardinia, starting around 28 Ma and peaking during the 22–18 Myr time range. This younger magmatic province undoubtedly has to be interpreted as related to W-directed Apenninic subduction and roll back. However, the Late Eocene magmatism at the Calabona locality was also interpreted to represent the effect of the northwest-directed subduction system that started to consume ancient oceanic lithosphere of the Tethys Ocean. This subduction would have initiated in a time span prior to the Late Eocene that is long enough to culminate in 100–150 km of subduction required to generate an arc (Lustrino et al. 2009). Given the very slow Africa–Europe convergence rates in the pre-Late Eocene (only 250 km of convergence occurred between Africa and Eurasia prior to 35 Ma at the position of Sardinia, and even less farther to the west; van Hinsbergen et al. 2014a), this would suggest that an Alpine-vergent thrusting in Sardinia and farther to the west would have been a back-thrust above a long-lived Apennines-type subduction. However later, Lustrino et al. (2013) argued, based on geochemical arguments, that the Eocene manifestations of the subduction related magmatism are due to the partial melting of Hercynian lower crust rather than to the dehydration processes of subducting Alpine Tethys oceanic crust. They pessimistically concluded that “whatever subduction dip, convergence rate and depth of partial melting are chosen, it is not possible to obtain a geologically sound model able to explain the spatial–temporal relationships of the western Mediterranean subduction-related igneous districts”.

Since the pioneering work of Boccaletti et al. (1971) another group of authors (e.g. Guerrero et al. 1993; Doglioni et al. 1999; Michard 2006; Molli 2008; Handy et al. 2010) favored the idea of a temporal rather than lateral change in subduction polarity for the entire Apennines area (see discussion in Michard et al. 2006, their Figure 9). The validity of this hypothesis is intimately linked to the question about kinematics and timing of thrusting, and, paleogeographic origin of the Calabrian–Peloritanean Belt that consists of high-grade Variscan basement nappes thrust on top of Ligurian-type ophiolites (Schenk 1990; Vitale and Ciarcia 2013). It is also related to the question as to whether the Calabrian belt is part of an ALKAPECA block (Bouillin et al. 1986; Michard et al. 2006; Handy et al. 2010) rather than a part of Sardinia, i.e. Europe.



The data and their interpretation presented in the previous chapters favor a temporal rather than lateral change in subduction polarity at least for the Northern Apennines and their transition into the Alps. However, a more detailed discussion of the southern Apennines of Calabria is outside the scope of this contribution. The controversy around a lateral vs. temporal change in subduction polarity in Calabria needs to be solved by future investigations in Calabria.

Although our data and interpretations support a temporal change in subduction polarity in the Northern Apennines realm and Corsica this interpretation faces a serious problem that is not yet solved. It is not easy to explain why this change occurred so shortly after the end of E-directed Alpine subduction in Corsica (about 35–37 Ma; Vitale Brovarone and Herwartz 2012). The switch in subduction polarity around Corsica is not precisely known but probably occurred in Latest Eocene to Early Oligocene times. According to kinematic and paleomagnetic reconstructions (S eranne 1999; Gattacceca et al. 2007; Advokaat et al. 2014) fast roll back of the East Ligurian oceanic crust and rotation of Corsica-Sardinia commenced at around 30 Ma, followed by rotation of Sardinia between 21 and 16 Ma. This only leaves some 5–7 Ma for slab break and change of subduction polarity and sufficient depth of subduction in order to trigger roll back of the W-dipping slab. Apenninic W-directed subduction and fast roll-back was certainly triggered by the negative buoyancy of large parts of the Piedmont-Liguria Ocean that still stayed open when Alpine orogeny in the Ligurian Alps ended at the end of the Eocene (Handy et al. 2010). This points to a change from plate convergence-driven Alpine E- to SE-directed subduction in Western Alps and Corsica toward W-directed subduction almost entirely driven by slab roll back in the Northern Apennines.

## 6 Summary and conclusions

The geological interpretation of the P-wave velocity model of Diehl et al. (2009) allows for depicting unprecedented details concerning the outlines of this Adriatic mantle wedge and integrates its outline into the overall structure of the Western Alps. Important along-strike changes are visualized in five radially arranged crustal-scale transects across the Alps (Fig. 2) and the tectonic maps of Figs. 1 and 6). The interpretation of these findings reveals an important difference between Eastern and Western Alps. While the upper plate of the Eastern Alps is defined by the Austroalpine nappe system consisting of crustal imbricates of continental material, the upper plate of the Western Alps is formed by non-metamorphic klippen of the Piedmont-Ligurian Oceanic crust and lithosphere (Molli 2008).

Based on this, and combined with a review of existing structural data, a 3-stage evolution is proposed for the formation of the arc of the Western Alps. A first stage (Late Cretaceous–35 Ma) is characterized by top-NNW thrusting in sinistral transpression causing at least some 260 km displacement of the E-W-striking Central and Eastern Alps, together with the Adria micro-plate, towards N to NNW with respect to stable Europe. A second stage (35–25 Ma) is characterized by lateral indentation of Adria including the Ivrea mantle wedge at its leading edge towards the WNW. This culminated in some 100–150 km of right-lateral strike-slip of the Ivrea mantle wedge, Western Alps and Adriatic micro-plate relative to the eastern Alps and Eurasia. This was accommodated in the Western Alpine nappe pile by top-WNW thrusting in the external zones of the central portion of the arc along the Penninic Frontal thrust combined with underthrusting of the Ivrea mantle wedge below the Western Alpine nappe pile. The third and final stage of arc formation (25–0 Ma) was associated with orogeny in the Apennines and led to oroclinal bending in the southernmost Western Alps, simultaneously with the 50° counterclockwise rotation of the Corsica-Sardinia block. Particularly during the second stage of the evolution of the arc of the Western Alps the Ivrea mantle wedge, exhumed to shallow crustal depth already during Jurassic rifting at the passive continental margin of the Adria continent and representing a high strength wedge, played a crucial role by controlling the lateral ends of indentation near Cuneo and the Rhone-Simplon fault.

The third stage of the evolution of the arc of the Western Alps calls for an analysis of the Alps–Apennines transition zone. A pre-requisite for understanding this transition is the realization that those parts of the Piedmont-Liguria oceanic domain that form the upper plate in the Western Alps remained open until the Late Eocene. A combination of the gravitational potential of a southwards enlarging domain of old and oceanic lithosphere, the arrival of the Corsican continent in the Alpine trench resisting subduction, and ongoing Africa-Europe convergence induced a change in subduction polarity from Alpine S to SE-directed subduction towards NW-directed subduction and roll back of the Apennines slab, pulled by the negative buoyancy of old oceanic lithosphere large parts of which remained unaffected by Alpine orogeny.

**Acknowledgements** We thank Peter Nievergelt for his great help during the preparation of the figures. The careful and critical reviews of Rob Butler and Niko Froitzheim helped much in improving an earlier version of the manuscript. We thank our colleagues of the new research initiative AlpArray for many inspiring discussions, and we look forward to learn about the new insights and concepts that will certainly result from this multi-disciplinary program.



## References

- Advokaat, E. L., van Hinsbergen, D. J. J., Maffione, M., Langereis, Cor G., Vissers, R. L. M., Cherchi, A., et al. (2014). Eocene rotation of Sardinia, and the paleogeography of the western Mediterranean region. *Earth and Planetary Science Letters*, *401*, 183–195.
- Argand, E. (1924). Des Alpes et de l’Afrique. *Bulletin de la Société vaudoise des sciences naturelles*, *55*, 233–236.
- Argnani, A. (2012). Plate motion and the evolution of Alpine Corsica and Northern Apennines. *Tectonophysics*, *579*, 207–219.
- Babist, J., Handy, M. R., Konrad-Schmolke, M., & Hammerschmidt, K. (2006). Precollisional, multistage exhumation of subducted continental crust: The Sesia Zone, western Alps. *Tectonics*, *25*, TC6008. doi:10.1029/2005TC001927.
- Barféty, J.C., Lemoine, M., de Graciansky, P.C., Tricart, P. & Mercier, D. (1995). *Notice explicative de la Carte géologique de France 1:50’000 Feuille Briançon (823)*. BRGM Orléans. 180 pp.
- Barnes, J., Beltrando, M., Lee, C. T., Cisneros, M., Loewy, S., & Chin, E. (2014). Geochemistry of Alpine serpentinites from rifting to subduction: A view across paleogeographic domains and metamorphic grade. *Chemical Geology*, *389*, 29–47.
- Bayer, R., Carozzo, M. T., Lanza, R., Miletto, M., & Rey, D. (1989). Gravity modeling along the ECORS-CROP vertical seismic reflection profile through the western Alps. *Tectonophysics*, *162*, 203–218.
- Beaumont, C., Fullsack, P. H., & Hamilton, J. (1994). Styles of crustal deformation in compressional orogens caused by subduction of the underlying lithosphere. *Tectonophysics*, *232*, 119–132.
- Bellahsen, N., Jolivet, L., Lacombe, O., Bellanger, M., Boutoux, A., Garcia, S., et al. (2012). Mechanisms of margin inversion in the external Western Alps: Implications for crustal rheology. *Tectonophysics*, *560–561*, 62–83.
- Beltrando, M., Compagnoni, R., Ferrando, S., Mohn, G., Frasca, G., Odasso, N. & Vukmanović, Z. (2015a). Adriatic margin: Canavese, Italy: Crustal thinning and mantle exhumation in the Levone area (Southern Canavese zone, Western Alps). *Journal of the Virtual Explorer*, *49*.
- Beltrando, M., Lister, G. S., Rosenbaum, G., Richards, S., & Forster, M. A. (2010). Recognizing episodic lithospheric thinning along a convergent plate margin: The example of the Early Oligocene Alps. *Earth-Science Reviews*, *103*, 81–98.
- Beltrando, M., Stöckli, D. F., Decarli, A., & Manatschal, G. (2015b). A crustal-scale view at rift localization along the fossil Adriatic margin of the Alpine Tethys preserved in NW Italy. *Tectonics*, *34*, 1927–1951.
- Berger, A. & Bousquet, R. (2008). Subduction-related metamorphism in the Alps: Review of isotopic ages based on petrology and their geodynamic consequences. In S. Siegesmund, B. Fügenschuh & N. Froitzheim (Eds.) *Tectonic ts of the Alpine-Dinaride-Carpathian System* (p. 117–144). Geological Society London, Special Publications, 298, 117–144.
- Berger, A. & Mercolli, I. (2006). Tectonic and Petrographic Map of the Central Lepontine Alps. *Carta geologica speciale 127, map sheet 43 Sopra Ceneri*. Federal Office of Topography swisstopo.
- Berger, A., Mercolli, I., & Engi, M. (2005). The central Lepontine Alps: Notes accompanying the tectonic and petrographic map sheet Sopra Ceneri (1:100’000). *Schweizerische Mineralogische und Petrographische Mitteilungen*, *85*, 109–146.
- Berger, A., Rosenberg, C., & Schmid, S. M. (1996). Ascent, emplacement and exhumation of the Bergell pluton within the Southern Steep Belt of the Central Alps. *Schweizerische Mineralogische und Petrographische Mitteilungen*, *76*, 357–382.
- Bernoulli, D., Bertotti, G., & Zingg, A. (1989). Northward thrusting of the Gonfolite Lombarda (“South-Alpine Molasse”) onto the Mesozoic sequence of the Lombardian Alps: Implications for the deformation history of the Southern Alps. *Eclogae Geologicae Helvetiae*, *82*, 841–856.
- Bernoulli, D., Giger, M., & Müller, D. W. (1993). Sr-isotope-geochronology of the Gonfolite Lombarda Group (“South-Alpine Molasse”, northern Italy) and radiometric constraints for its age of deposition. *Eclogae Geologicae Helvetiae*, *86*, 751–767.
- Bernoulli, D., Manatschal, G., Desmurs, L., and Müntener, O. (2003). Where did Gustav Steinmann see the trinity? Back to the roots of an Alpine ophiolite concept. In Y. Dilek, & S. Newcomb, S. (Eds.), *Ophiolite concept and the evolution of geological thought* (pp. 93–110). Boulder, Colorado: Geological Society of America Special Paper 373.
- Bertotti, G. (1991). Early Mesozoic extension and alpine shortening in the western Southern Alps: The geology of the area between Lugano and Menaggio (Lombardy, Northern Italy). *Memorie di Scienze Geologiche Padova*, *XLIII*, 17–123.
- Bertotti, G., Picotti, V., Bemoulli, D., & Castellarin, A. (1993). From rifting to drifting: Tectonic evolution of the South-Alpine upper crust from the Triassic to the Early Cretaceous. *Sedimentary Geology*, *86*, 53–76.
- Blundell, D. J., Freeman, R., & Mueller, S. (1992). *A continent revealed. The European Geotraverse, structure and dynamic evolution*. Cambridge: Cambridge University Press.
- Boccaletti, M., Elter, P., & Guazzone, G. (1971). Plate tectonic models for the development of the Western Alps and Northern Apennines. *Nature Physical Science*, *234*(49), 108–111.
- Bouillin, J.-P., Durand-Delga, M., & Olivier, P. (1986). Betic-Rifian and Tyrrhenian Arcs: Distinctive features, genesis, and development stages. In F. C. Wezel (Ed.), *The origin of arcs* (pp. 281–304). Amsterdam: Elsevier.
- Bousquet, R., Goffé, B., Henry, P., LePichon, X., & Chopin, C. (1997). Kinematic, thermal and petrological model of the Central Alps: Lepontine metamorphism in the upper crust and eclogitization of the lower crust. *Tectonophysics*, *273*, 105–127.
- Bousquet, R., Goffé, B., Vidal, O., Oberhänsli, R., & Patriat, M. (2002). The tectono-metamorphic history of the Valaisan domain from the Western to the Central Alps: New constraints for the evolution of the Alps. *Geological Society of America Bulletin*, *114*, 207–225.
- Bousquet, R., Oberhänsli, R., Schmid, S.M., Berger, A., Wiederkehr, M., Robert, C., Möller, A., Rosenberg, C., Zeilinger, G., Molli, G. & Koller, F. (2012a). Metamorphic framework of the Alps. Commission for the Geological Map of the World; Subcommission for Magmatic and Metamorphic Maps. IUGS and IUGG, Paris. <http://www.ccgmg.org>.
- Bousquet R., Schmid S.M., Zeilinger G, Oberhänsli R., Rosenberg C., Molli G., Robert C., Wiederkehr M., Rossi Ph. (2012b). *Tectonic framework of the Alps. Map 1: 1 000 000*. Commission for the Geological Map of the World; Subcommission for Magmatic and Metamorphic Maps. IUGS and IUGG, Paris. <http://www.ccgmg.org>.
- Boyden, J.A., Müller, R.D., Gurnis, M., Torsvik, T.H., Clark, J.A., Turner, M., Ivey-Law, H., Watson, R.J. & Cannon, J.S. (2011). Next-generation plate-tectonic reconstructions using GPLates. In G.R. Keller & C. Baru (Eds), *Geoinformatics* (pp. 95–114). Cambridge: Cambridge University Press.
- Brunet, C., Monié, P., Jolivet, L., & Cadet, J. P. (2000). Migration of compression and extension in the Tyrrhenian Sea, insights from 40Ar/39Ar ages on micas along a transect from Corsica to Tuscany. *Tectonophysics*, *321*, 127–155.
- Bucher, S., Schmid, S. M., Bousquet, R., & Fügenschuh, B. (2003). Late-stage deformation in a collisional orogen (Western Alps):

- nappe refolding, back-thrusting or normal faulting? *Terra Nova*, 15, 109–117.
- Bucher, S., Ulardic, C., Bousquet, R., Ceriani, S., Fügenschuh, B., & Schmid, S. M. (2004). Tectonic evolution of the Briançonnais units along the ECORS-CROP transect through the Italian-French Western Alps. *Eclogae Geologicae Helveticae*, 97, 321–345.
- Bürgisser, J., & Ford, M. (1998). Overthrust shear deformation of a foreland basin; structural studies south-east of the Pelvoux massif, SE France. *Journal of Structural Geology*, 20, 1455–1475.
- Burkhard, M. (1988). L'Helvétique de la bordure occidentale du massif de l'Aar (évolution tectonique et métamorphique). *Eclogae Geologicae Helveticae*, 81, 63–114.
- Burkhard, M. & Sommaruga A. (1998). Evolution of the western Swiss Molasse basin: Structural relations with the Alps and the Jura belt. In: Mascle C. et al. (Eds): Cenozoic Foreland Basins of Western Europe. *Special Publications Geological Society London*, 134, 279–298.
- Butler, R. W. H. (1983). Balanced cross sections and their implications for the deep structure of the northwest Alps. *Journal of Structural Geology*, 5, 125–137.
- Butler, R. W. H. (1986). Thrust tectonics, deep structure and crustal subduction in the Alps and Himalayas. *Journal of the Geological Society of London*, 143, 857–873.
- Butler, R. W. H. (1992). Thrust zone kinematics in a basement-cover imbricate stack; eastern Pelvoux Massif, French Alps. *Journal of Structural Geology*, 14, 29–40.
- Butler, R. H. W. (2013). Area balancing as a test of models for the deep structure of mountain belts, with specific reference to the Alps. *Journal of Structural Geology*, 52, 2–16.
- Byerlee, J. (1978). Friction of rocks. *Pure and Applied Geophysics*, 116, 615–626.
- Caby, R. (1996). Low-angle extrusion of high-pressure rocks and the balance between outward and inward displacements of Middle Penninic units in the western Alps. *Eclogae Geologicae Helveticae*, 89(1), 229–267.
- Capponi, G., Crispini, L. & Scambelluri, M. (2009). Discussion: Comment on “Subduction polarity reversal at the junction between the Western Alps and the Northern Apennines, Italy”, by G. Vignaroli, C. Faccenna, L. Jolivet, C. Piromallo, F. Rossetti. *Tectonophysics*, 465, 221–226.
- Cardello, G. L., & Mancktelow, N. (2015). Veining and post-nappe transensional faulting in the SW Helvetic Alps (Switzerland). *Swiss Journal of Geosciences*, 108, 379–400.
- Carey, S. W. (1955). The orocline concept in geotectonics, part I. *Papers and Proceedings of the Royal Society of Tasmania*, 89, 255–288.
- Carminati, E., Doglioni, C. & Scrocca, D. (2004). Alps vs Apennines. In U. Crescenti *Geology of Italy: Special Volume of the Italian Geological Society for the IGC 32 Florence 2004* (pp. 141–151). Roma: Società Geologica Italiana.
- Carrapa, B., & Garcia-Castellanos, D. (2005). Western Alpine back-thrusting as subsidence mechanism in the Tertiary Piedmont Basin (Western Po Plain, NW Italy). *Tectonophysics*, 406, 197–212.
- Cassano, E., Anelli, L., Fichera, R. & Cappelli, V. (1986). *Pianura Padana. Interpretazione integrate di dati geofisici e geologici*. 73. Congresso Società Geologica Italiana. Centro Stampa AGIP Roma, pp. 1–28.
- Castellarin, A. (2001). Alps–Apennines and Po–Plain—Frontal Apennines relationships. In G.B. Vai et al., *Anatomy of an Orogen. The Apennines and adjacent Mediterranean Basins* (pp. 177–196). London: Kluwer.
- Ceriani, S., Fügenschuh, B., & Schmid, S. M. (2001). Multi-stage thrusting at the “Penninic Front” in the Western Alps between Mont Blanc and Pelvoux massifs. *International Journal of Earth Sciences*, 90, 685–702.
- Ceriani, S., & Schmid, S. M. (2004). From N-S collision to WNW-directed post-collisional thrusting and folding: Structural study of the Frontal Penninic Units in Savoie (Western Alps, France). *Eclogae Geologicae Helveticae*, 97, 347–369.
- Chemenda, A. L., Mattauer, M., Malavieille, J., & Bokun, A. N. (1995). A mechanism for syn-collisional rock exhumation and associated normal faulting: Results from physical modeling. *Earth and Planetary Science Letters*, 132, 225–232.
- Collombet, M., Thomas, J. C., Chauvin, A., Tricart, P., Bouillin, J. P., & Gratier, J. P. (2002). Counterclockwise rotation of the western Alps since the Oligocene: New insights from paleomagnetic data. *Tectonics*. doi:10.1029/2001TC901016.
- Compagnoni, R., Dal Piaz, G. V., Hunziker, J. C., Gosso, G., Lombardo, B., & Williams, P. F. (1977). The Sesia-Lanzo Zone, a slice of continental crust with Alpine high pressure-low temperature assemblages in the Western Italian Alps. *Rendiconti della Società Italiana di Mineralogia e Petrologia*, 33, 281–334.
- Cortesogno, L., & Haccard, D. (1984). Note illustrative alla carta geologica della Zona Sestri-Voltaggio. *Memorie Società Geologica Italiana*, 28, 115–150.
- Crispini, L., Federico, L., Cappini, G., & Spagnolo, C. (2009). Late orogenic transpressional tectonics in the «Ligurian Knot». *Italian Journal of Geosciences*, 128, 433–441.
- Dal Piaz, G. V., Cortiana, G., Del Moro, A., Martin, S., Pennacchioni, G., & Tartarotti, P. (2001). Tertiary age and paleostructural inferences of the eclogitic imprint in the Austroalpine outliers and Zermatt-Saas ophiolite, western Alps. *International Journal of Earth Sciences*, 90, 668–684.
- de Graciansky, P. C., Dardeau, G., Lemoine, M., & Tricart, P. (1989). The inverted margin of the French Alps and foreland basin inversion. *Geological Society London Special Publications*, 44, 87–104.
- Decandia, F. A., & Elter, P. (1972). La ‘zona’ ofiolitifera del Bracco nel settore compreso tra Levante e la Val Graveglia (Appennino ligure). *Memorie della Società Geologica Italiana*, 11, 503–530.
- Delacou, B., Sue, C., Champagnac, J. D., & Burkhard, M. (2004). Present-day geodynamics in the bend of the western and central Alps as constrained by earthquake analysis. *Geophysical Journal International*, 158, 753–774.
- Di Giulio, A. (1999). Mass transfer from the Alps to the Apennines: volumetric constraints in the provenance study of the Macigno-Modino source–basin system, Chatian-Aquitania, northwestern Italy. *Sedimentary Geology*, 124, 69–80.
- Di Giulio, A. & Galbiati, B. (1998). Interaction between tectonics and deposition into an episutural basin in the Alps-Apennine knot. In R. Polino & R. Sacchi (Eds.), *Atti del convegno “Rapporti Alpi-Appennino”* (pp. 113–128). Accademia Nazionale delle Scienze detta dei XL, 14, Roma.
- Diehl, T., Husen, S., Kissling, E., & Deichmann, N. (2009). High-resolution 3-D P-wave model of the Alpine crust. *Geophysical Journal International*, 179, 1133–1147.
- Doglioni, C., Fernandez, M., Gueguen, E., & Sabat, F. (1999). On the interference between the early Apennines-Maghrebides back-arc extension and the Alps-Betics orogen in the Neogene geodynamics of the Western Mediterranean. *Bollettino Società Geologica Italiana*, 118, 75–89.
- Doglioni, C., Mongelli, F., & Pialli, G. (1998). Boudinage of the Alpine belt in the Apenninic back-arc. *Memorie della Società Geologica Italiana*, 52, 457–468.
- Druiventak, A., Trepmann, C. A., Renner, J., & Hanke, K. (2011). Low-temperature plasticity of olivine during high stress deformation of peridotite at lithospheric conditions—an experimental study. *Earth and Planetary Science Letters*, 311, 199–211.

- Dumont, T., Champagnac, J. D., Crouzet, C., & Rochat, P. (2008). Multistage shortening in the Dauphiné zone (French Alps): The record of Alpine collision and implications for pre-Alpine restoration. *Swiss Journal of Geosciences*, *101*(Supplement 1), 89–110.
- Dumont, T., Simon-Labric, T., Authemayou, C., & Heymes, T. (2011). Lateral termination of the north-directed Alpine orogeny and onset of westward escape in the Western Alpine arc: Structural and sedimentary evidence from the external zone. *Tectonics*. doi:10.1029/2010TC002836.
- Dumont, T., Schwartz, S., Guillot, S., Simon-Labric, T., Tricart, P., & Jourdan, S. (2012). Structural and sedimentary records of the Oligocene revolution in the Western Alpine arc. *Journal of Geodynamics*, *56–57*, 18–28.
- Durand-Delga, M. (1984). Principaux traits de la Corse Alpine et corrélations avec les Alpes Ligures. *Memorie della Società Geologica Italiana*, *28*, 285–329.
- Ellero, A., Leoni, L., Marroni, M., & Sartori, F. (2001). Internal Liguride Units from Central Liguria, Italy: new constraints to the tectonic setting from white mica and chlorite studies. *Schweizerische Mineralogische und Petrographische Mitteilungen*, *81*, 39–53.
- Elter, G., Elter, P., Sturani, C., & Weidmann, M. (1966). Sur la prolongation du domaine ligure de l'Apennin dans le Montferrat et les Alpes et sur l'origine de la Nappe de Simme s.l. des Préalpes romandes et chablaisiennes. *Archives des Sciences*, *19*, 279–377.
- Elter, P., & Pertusati, P. (1973). Considerazioni sul limite Alpi-Apennino a sulle sue relazioni con l'arco delle Alpi Occidentali. *Memorie della Società Geologica Italiana*, *12*, 359–375.
- Escher, A., Hunziker, J.C., Marthaler, M., Masson, H., Sartori, M., & Steck, A. (1997). Geologic framework and structural evolution of the western Swiss-Italian Alps. In: Pfiffner A.O. et al. (Eds.): *Deep structure of the Swiss Alps: Results from NRP 20* (pp. 205–221). Basel, Switzerland: Birkhäuser.
- Faccenna, C., Jolivet, L., Piromallo, C., & Morelli, A. (2003). Subduction and the depth of convection in the Mediterranean mantle. *Journal of Geophysical Research*. doi:10.1029/2001JB001690.
- Fassmer, K., Obermüller, G., Nagel, T. J., Kirst, F., Froitzheim, N., Sandmann, S., et al. (2016). High-pressure metamorphic age and significance of eclogite-facies continental fragments associated with oceanic lithosphere in the Western Alps (Etirol-Levaz Slice, Valtourmenche, Italy). *Lithos*, *252–253*, 145–159.
- Festa, A., & Codegone, G. (2013). Geological map of the External Ligurian Units in western Monferrato (Tertiary Piedmont Basin, NW Italy). *Journal of Maps*, *9*, 84–97.
- Finetti, I. R. (Ed.). (2005). *CROP PROJECT: Deep Seismic Exploration of the Central Mediterranean and Italy (779 pp & 72 plates)*. Amsterdam: Elsevier.
- Finetti, I. R., Boccaletti, M., Bonini, M., Del Ben, A., Geletti, R., Pipam, N., et al. (2001). Crustal section based on CROP seismic data across the North Tyrrhenian-Northern Apennines–Adriatic Sea. *Tectonophysics*, *343*, 135–163.
- Ford, M., Duchêne, S., Gasquet, D., & Vanderhaeghe, O. (2006). Two-phase orogenic convergence in the external and internal SW Alps. *Journal of the Geological Society, London*, *163*, 815–826.
- Ford, M. & Lickorish, W.H. (2004). Foreland basin evolution around the western Alpine Arc. In: P. Joseph & S.A. Lomas (Eds.), *Deep-water sedimentation in the Alpine Basin of SE France: New perspectives on the Grès d'Annot and related systems* (pp. 39–63). Geological Society London Special Publication 221.
- Fountain, D. M. (1976). The Ivrea-Verbanò and Strona-Ceneri zones, Northern Italy: A cross-section of the continental crust: new evidence from seismic velocities of rock samples. *Tectonophysics*, *33*, 145–165.
- Fox, M., Herman, F., Willett, S. D., & Schmid, S. M. (2016). The exhumation history of the European Alps inferred from linear inversion of thermochronometric data. *American Journal of Science*, *316*, 505–541.
- Froitzheim, N., Schmid, S. M., & Frey, M. (1996). Mesozoic paleogeography and the timing of eclogite-facies metamorphism in the Alps: A working hypothesis. *Eclogae Geologicae Helvetiae*, *89*, 81–110.
- Fry, N. (1989). Southwestward thrusting and tectonics of the western Alps. In: M.P. Coward et al. (Eds.), *Alpine Tectonics* (pp. 83–109). Geological Society London Special Publication 45.
- Fügenshuh, B., & Schmid, S. M. (2003). Late stages of deformation and exhumation of an orogen constrained by fission track data: A case study in the Western Alps. *Geological Society of America Bulletin*, *115*, 1425–1440.
- Fujimoto, Y., Konob, Y., Hirajima, T., Kanagawa, K., Ishikawa, M., & Arima, M. (2010). P-wave velocity and anisotropy of lawsonite and epidote blueschists: Constraints on water transportation along subducting oceanic crust. *Physics of the Earth and Planetary Interiors*, *183*, 219–228.
- Ganne, J., Marquer, D., Rosenbaum, G., Bertrand, J. M., & Fudral, S. (2006). Partitioning of deformation within a subduction channel during exhumation of high-pressure rocks: A case study from the Western Alps. *Journal of Structural Geology*, *28*, 1193–1207.
- Garzanti, E., & Malusa, M. G. (2008). The Oligocene Alps: Domal unroofing and drainage development during early orogenic growth. *Earth and Planetary Science Letters*, *268*, 487–500.
- Gattacceca, J., Deino, A., Rizzo, R., Jones, D. S., Henry, B., Beaudoin, B., et al. (2007). Miocene rotation of Sardinia: New paleomagnetic and geochronological constraints and geodynamic implications. *Earth and Planetary Science Letters*, *258*, 359–377.
- Gebauer, D., Schertl, H. P., Brix, M., & Schreyer, W. (1997). 35 Ma old ultrahigh-pressure metamorphism and evidence for very rapid exhumation in the Dora Maira Massif, Western Alps. *Lithos*, *41*, 5–24.
- Gidon, M. (1972). Les chaînons briançonnais et subbriançonnais de la rive gauche de la Stura entre le Val de l'Arma (province de Cuneo Italie). *Géologie Alpine*, *48*, 87–120.
- Giuntoli, F. & Engi, M. (2016). Internal geometry of the central Sesia Zone (Aosta Valley, Italy): HP tectonic assembly of continental slices. *Swiss Journal of Geosciences*. doi:10.1007/s00015-016-0225-4.
- Grandjacquet, C., & Haccard, D. (1977). Position structural et rôle paléogéographique de l'unité du Bracco au sein du context ophiolitique liguro-piémontais (Apennin—Italie). *Bulletin de la Société géologique de France*, *19*, 901–904.
- Grasemann, B., & Mancktelow, N. S. (1993). Two-dimensional thermal modeling of normal faulting: The Simplon fault zone, central Alps, Switzerland. *Tectonophysics*, *225*, 155–165.
- Guerrera, F., Martín-Algarra, A., & Perrone, V. (1993). Late Oligocene-Miocene syn-/late-orogenic successions in Western and Central Mediterranean Chains from the Betic Cordillera to the Southern Apennines. *Terra Nova*, *5*, 525–544.
- Guillot, S., & Ménot, R. P. (2009). Paleozoic evolution of the External Crystalline Massifs of the Western Alps. *Comptes Rendus Geoscience*, *341*, 253–265.
- Handy, M. R. (1987). The structure, age and kinematics of the Pogallo Fault Zone; Southern Alps, northwestern Italy. *Eclogae Geologicae Helvetiae*, *80*, 593–632.
- Handy, M. R., Franz, L., Heller, F., Janott, B., & Zurbriggen, R. (1999). Multistage accretion and exhumation of the continental crust (Ivrea crustal section, Italy and Switzerland). *Tectonics*, *18*, 1154–1177.



- Handy, M. R., Schmid, S. M., Bousquet, R., Kissling, E., & Bernoulli, D. (2010). Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological–geophysical record of spreading and subduction in the Alps. *Earth-Science Reviews*, *102*, 121–158.
- Handy, M. R., & Zingg, A. (1991). The tectonic and rheologic evolution of an attenuated cross-section of the continental crust: Ivrea crustal section, Southern Alps, northwestern Italy and southern Switzerland. *Geological Society of America Bulletin*, *103*, 236–253.
- Henry, C., Michard, A., & Chopin, C. (1993). Geometry and structural evolution of ultra-high-pressure and high-pressure rocks from the Dora-Maira massif, Western Alps. *Journal of Structural Geology*, *15*(8), 965–981.
- Huber, M., Ramsay, J., & Simpson, C. (1980). Deformation of the Maggia and Antigorio nappes Lepontine Alps. *Eclogae Geologicae Helveticae*, *73*, 593–606.
- Jaillard, E. (1999). The Late Cretaceous-Eocene sedimentation in the internal Briançonnais units of Vanoise (French Alps): Witnesses of early Alpine movements. *Eclogae Geologicae Helveticae*, *92*, 211–220.
- Jolivet, L., Faccenna, C., Goffé, B., Mattei, M., Rossetti, F., Brunet, C., et al. (1998). Mid-crustal shear zones in postorogenic extension: Example from the northern Tyrrhenian Sea. *Journal of Geophysical Research*, *103*, 12123–12160.
- Keller, L., Fügenschuh, B., Hess, M., Schneider, B., & Schmid, S. M. (2006). Simplon fault zone in the western and central Alps: Mechanism of Neogene faulting and folding revisited. *Geology*, *34*, 317–320. doi:10.1130/G22256.1.
- Keller, L. M., & Schmid, S. M. (2001). On the kinematics of shearing near the top of the Monte Rosa nappe and the nature of the Furgg zone in Val Loranco (Antrona valley, N. Italy): Tectonometamorphic and paleogeographical consequences. *Schweizerische Mineralogische und Petrographische Mitteilungen*, *81*, 347–367.
- Kerckhove, C. (1969). La “zone du Flysch” dans les nappes de l’Embrunais-Ubaye (Alpes occidentales). *Géologie Alpine*, *45*, 5–204.
- Kissling, E. (1984). Three-dimensional gravity model of the Northern Ivrea-Verbano zone. In Wagner, J.J. & Mueller, S. (Eds.): *Matériaux pour la Géologie de la Suisse Géophysique No 21. Geomagnetic and gravimetric studies of the Ivrea Zone*. Bern: Kümmerly & Frey, Geographischer Verlag, 1–61.
- Kramer, J. (2002). *Structural evolution of the Penninic units in the Monte Rosa region (Swiss and Italian Alps)* (144 pp.). Ph.D. Basel University, edoc University Library of Basel.
- Lacombe, O., & Jolivet, L. (2005). Structural and kinematic relationships between Corsica and the Pyrenees-Provence domain at the time of the Pyrenean orogeny. *Tectonics*, *24*, TC1003. doi:10.1029/2004TC001673.
- Lahondère, D. (1996). *Les schistes bleus et les éclogites à lawsonite des unités continentales et océanique de la Corse alpine: Nouvelles données pétrologiques et structurales (Corse)*. France: BRGM.
- Lardeaux, J. M., Schwartz, S., Tricart, P., Paul, A., Guillot, S., Béthoux, N., et al. (2006). A crustal-scale cross-section of the south-western Alps combining geophysical and geological imagery. *Terra Nova*, *18*(6), 412–422.
- Laubscher, H. P. (1971). The large-scale kinematics of the Western Alps and the Northern Apennines and its palinspastic implications. *American Journal of Science*, *271*, 193–226.
- Laubscher, H. (1991). The arc of the Western Alps today. *Eclogae Geologicae Helveticae*, *84*, 631–659.
- Laubscher, H., Biella, G. C., Cassinis, R., Gelati, R., Lozej, A., Scarascia, S., et al. (1992). The collisional knot in Liguria. *Geologische Rundschau*, *81*(2), 275–289.
- Lemoine, M., & Tricart, P. (1986). Les Schistes lustrés piémontais des Alpes Occidentales: Approche stratigraphique, structural et sédimentologique. *Eclogae Geologicae Helveticae*, *79*(2), 271–294.
- Leu, W. (1987). Lithostratigraphie und Tektonik der nordpenninischen Sedimente in der Region Bedretto-Baceno-Visp. *Eclogae Geologicae Helveticae*, *79*, 769–824.
- Levi, N., Ellero, A., Ottria, G., & Pandolfi, L. (2006). Polyorogenic deformation history recognized at very shallow structural levels: The case of the Antola Unit (Northern Apennine, Italy). *Journal of Structural Geology*, *28*, 1694–1709.
- Lickorish, W.H. & Ford, M. (1998). Evolution of the Digne thrust system, southern Subalpine chains: kinematics and timing of deformation. In A. Mascle et al. (Eds.), *Cenozoic Foreland Basins of Western Europe* (pp. 189–211). Geological Society London, Special Publications 134.
- Lippitsch, R., Kissling, E., & Ansorge, J. (2003). Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography. *Journal of Geophysical Research*, *108*(B8), 2376. doi:10.1029/2002JB002016.
- Loprieno, A., Bousquet, R., Bucher, S., Ceriani, S., Dalla Torre, F. H., Fügenschuh, B., et al. (2011). The Valais units in Savoy (France): A key area for understanding the paleogeography and the tectonic evolution of the Western Alps. *International Journal of Earth Sciences*, *100*, 963–992.
- Lustrino, M., Fedele, L., Melluso, L., Morra, V., Ronga, F., Geldmacher, J., et al. (2013). Origin and evolution of Cenozoic magmatism of Sardinia (Italy). A combined isotopic (Sr–Nd–Pb–O–Hf–Os) and petrological view. *Lithos*, *180–181*, 138–158.
- Lustrino, M., Morra, V., Fedele, L., & Franciosi, L. (2009). Beginning of the Apennine subduction system in central western Mediterranean: Constraints from Cenozoic “orogenic” magmatic activity of Sardinia, Italy. *Tectonics*, *28*, TC5016. doi:10.1029/2008TC002419.
- Maffione, M., Speranza, F., Faccenna, C., Cascella, A., Vignaroli, G., & Sagnotti, L. (2008). A synchronous Alpine and Corsica-Sardinia rotation. *Journal of Geophysical Research*, *113*, B03104. doi:10.1029/2007JB005214.
- Malaroda, R., Carraro, F., Dal Piaz, G.V., Franceschetti, B., Sturani, C. & Zanella, E. (1970). Carta geologica del Massiccio dell’Argentera alla scala 1:50.000 e Note illustrative. *Memorie della Società Geologica Italiana*, *IX*, 557–663.
- Malavieille, J., Chemenda, A., & Larroque, C. (1998). Evolutionary model for Alpine Corsica: Mechanism for ophiolite emplacement and exhumation of high-pressure rocks. *Terra Nova*, *10*, 317–322.
- Malavieille, J., Lacassin, R., & Mattauer, M. (1984). Signification tectonique des lineations d’allongement dans les Alpes occidentales. *Bulletin de la Société Géologique de France*, *26*(5), 895–906.
- Manatschal, G., Sauter, D., Karpoff, A. M., Masini, E., Mohn, G., & Lagabriele, Y. (2011). The Chenaillet Ophiolite in the French/Italian Alps: An ancient analogue for an Oceanic Core Complex? *Lithos*, *124*, 169–184.
- Mancktelow, N. S. (1992). Neogene lateral extension during convergence in the Central Alps: Evidence from interrelated faulting and backfolding around the Simplonpass (Switzerland). *Tectonophysics*, *215*, 295–317.
- Manzotti, P., Ballèvre, M., Zucchi, M., Robyr, M., & Engi, M. (2014). The tectonometamorphic evolution of the Sesia-Dent Blanche nappes (internal Western Alps): review and synthesis. *Swiss Journal of Geosciences*, *107*, 309–336.
- Marchant, R. (1993). The Underground of the Western Alps. *Mémoires de Géologie (Lausanne)*, *15*, 1–137.
- Marchant, R. & Stampfli, G.M. (1997). Geodynamic evolution of the Tethyan margins of the Western Alps: Geodynamic significance. In O.A. Pfiffner, P. Lehner, P. Heitzmann, S. Mueller, A. &



- Steck (Eds.). *Deep structure of the Swiss Alps—results from NRP 20* (pp. 139–153). Basel: Birkhäuser.
- Marroni, M., Cerrina Feroni, A., di Biase, D., Ottria, G., Pandolfi, L., & Taini, A. (2002a). Polyphase folding at upper structural levels in the Borbera Valley (northern Apennines, Italy): Implications for the tectonic evolution of the linkage area between Alps and Apennines. *Comptes Rendus Geoscience*, *334*, 565–572.
- Marroni, M., Meneghini, F., & Pandolfi, L. (2010). Anatomy of the Ligure-Piemontese subduction system: Evidence from Late Cretaceous–middle Eocene convergent margin deposits in the Northern Apennines, Italy. *International Geology Review*, *52*, 1160–1192.
- Marroni, M., Molli, G., Montanini, A., Ottria, G., Pandolfi, L., & Tribuzio, R. (2002b). The External Ligurian units (Northern Apennine, Italy): From rifting to convergence of a fossil ocean-continent transition zone. *Ophioliti*, *27*, 119–131.
- Marroni, M., Molli, G., Ottria, G., & Pandolfi, L. (2001). Tectono-sedimentary evolution of the External Liguride units (Northern Apennines, Italy): Insights in the pre-collisional history of a fossil ocean-continent transition zone. *Geodinamica Acta*, *14*, 307–320.
- Marroni, M., Molli, G., Pandolfi, L., & Taini, A. (1999). Foliated cataclasites at the base of the Antola unit (Italy): Structural features and geological implications. *Comptes de l'Académie des Sciences Series IIA, Earth and Planetary Science*, *329*, 135–141.
- Marroni, M., & Pandolfi, L. (1996). The deformation history of an accreted ophiolite sequence: the Internal Liguride units (Northern Apennines, Italy). *Geodinamica Acta*, *9*, 13–29.
- Marroni, M., & Pandolfi, L. (2003). Deformation history of the ophiolite sequence from the Balagne Nappe, northern Corsica: Insights in the tectonic evolution of Alpine Corsica. *Geological Journal*, *38*, 67–83.
- Marroni, M., Molli, G., Montanini, A., & Tribuzio, R. (1998). The association of continental crust rocks with ophiolites in the Northern Apennines (Italy): implications for continent-ocean transition in the Western Tethys. *Tectonophysics*, *292*, 43–66.
- Marshak, S. (1988). Kinematics of orocline and arc formation in thin-skinned orogens. *Tectonics*, *7*, 73–86.
- Marton, E., Zampieri, D., Kazmer, M., Dunkl, I., & Frisch, W. (2011). New Paleocene-Eocene paleomagnetic results from the foreland of the Southern Alps confirm decoupling of stable Adria from the African plate. *Tectonophysics*, *504*, 89–99.
- Mattauer, M., Faure, M., & Malavieille, J. (1981). Transverse lineation and large-scale structures related to Alpine obduction in Corsica. *Journal of Structural Geology*, *3*, 401–409.
- Mauffret, A., Contrucci, I., & Brunet, C. (1999). Structural evolution of the Northern Tyrrhenian Sea from new seismic data. *Marine and Petroleum Geology*, *16*, 381–407.
- Ménard, G., & Molnar, P. (1991). Budget of crustal shortening and subduction of continental crust in the Alps. *Tectonics*, *10*, 231–244.
- Menkveld, J.W. (1995). *Der geologische Bau des Helvetikums der Innerschweiz* (165 pp & 141 attachments). Dissertation Universität Bern.
- Merle, O., & Brun, J. P. (1984). The curved translation path of the Parpaillon Nappe (French Alps). *Journal of Structural Geology*, *6*(6), 7111–7719.
- Michard, A., Avigad, D., Goffé, B., & Chopin, C. (2004). The high-pressure metamorphic front of the south Western Alps (Ubaye-Maira transect, France). *Schweizerische Mineralogische und Petrographische Mitteilungen*, *84*, 215–235.
- Michard, A., Negro, F., Saddiqi, O., Bouybaouene, M. L., Chalouan, A., Montigny, R., et al. (2006). Pressure–temperature–time constraints on the Maghrebide mountain building: evidence from the Rif-Betic transect (Marocco, Spain), Algerian correlations, and geodynamic implications. *Comptes Rendus Geoscience*, *338*, 92–114.
- Milnes, A. G. (1974). Structure of the Pennine Zone (Central Alps): A new working hypothesis. *Geological Society of America Bulletin*, *85*, 1727–1732.
- Mohn, G., Manatschal, G., Beltrando, M., & Hauptert, I. (2014). The role of rift-inherited hyper-extension in Alpine-type orogens. *Terra Nova*, *26*, 347–353.
- Molli, G. (1996). Pre-orogenic tectonic framework of the northern Apennine ophiolites. *Eclogae Geologicae Helveticae*, *89*, 163–180.
- Molli, G. (2008). Northern Apennine–Corsica orogenic system: an updated overview. In S. Siegesmund et al. (Eds.) *Tectonic aspects of the Alpine-Dinaride-Carpathian system* (pp. 413–442). Geological Society London Special Publications, 298.
- Molli, G., Crispini, L., Mosca, P., Piana, P. & Federico, L. (2010). Geology of the Western Alps-Northern Apennine junction area: a regional review. *Journal of the Virtual Explorer* *36*, paper 10 of the electronic edition ISSN 1441-8142.
- Molli, G., & Malavieille, J. (2011). Orogenic processes and the Corsica/Apennines geodynamic evolution: Insights from Taiwan. *International Journal of Earth Sciences*, *100*, 1207–1224.
- Molli, G., Tribuzio, R., & Marquer, D. (2006). Deformation and metamorphism at the eastern border of the Tenda Massif (NE Corsica): A record of subduction and exhumation of continental crust. *Journal of Structural Geology*, *28*, 1748–1766.
- Mosca, P., Polino, R., Rogledi, S., & Rossi, M. (2010). New data for the kinematic interpretation of the Alps–Apennines junction (Northwestern Italy). *International Journal of Earth Sciences*, *99*, 833–849.
- Müller, F. (1938). Geologie der Engelhörner, der Aareschlucht und der Kalkkeile bei Innertkirchen (Berner Oberland). *Beiträge zur Geologischen Karte der Schweiz, NF 74*, 71 pp.
- Mutti, E., Papani, L., Di Biase, D., Davoli, G., Mora, S., Segadelli, S., et al. (1995). Il Bacino Terziario Epimesoalpino e le sue implicazioni sui rapporti tra Alpi ed Appennino. *Memorie di Scienze Geologiche di Padova*, *47*, 217–244.
- Oberhänsli, R. (Ed) (2004). Metamorphic structure of the Alps. *Commission for the Geological Map of the World; Subcommission for Magmatic and Metamorphic Maps*. Paris: IUGS and IUGG. <http://www.ccg.org>.
- Orti, L., Pandeli, E., & Principi, G. (2002). New geological data from Gorgona Island (Northern Tyrrhenian Sea). *Ophioliti*, *27*, 133–144.
- Padovano, M., Elter, F. M., Pandeli, E., & Franceschelli, M. (2012). The East Variscan Shear Zone: New insights into its role in the Late Carboniferous collision in southern Europe. *International Geology Review*, *54*, 957–970.
- Pertusati, P. C., & Horrenberger, J. C. (1975). Studio strutturale degli scisti di Val Lavagna (Unità del Monte Gottero, Apennino Ligure). *Bollettino Società Geologica Italiana*, *94*, 1375–1436.
- Pfiffner, O.A. (2011). Cross sections of the Helvetic nappe system Plates V & IV. In: Pfiffner et al. 2011: *Structural map of the Helvetic Zone of the Swiss Alps. Geological Special Map 1:100'000*. Federal Office of Topography swisstopo.
- Pfiffner, O.A., Hänni, R., Kammer, A., Kligfield, R., Menkveld, J.W., Ramsay, J.G., Schmid, S.M. & Zurriggen, R. (2010). Geological Special Map 128/4 Sheet 37 Brünigpass. In: Pfiffner et al. 2011: *Structural map of the Helvetic Zone of the Swiss Alps. Geological Special Map 1:100'000*. Federal Office of Topography swisstopo.
- Pfiffner, O. A., & Heitzmann, P. (1997). Geological interpretation of the seismic profiles of the Central Traverse (lines C1, C2 and C3-north). In A. O. Pfiffner et al. (Eds.), *Deep structure of the Swiss Alps: Results from NRP 20* (pp. 115–122). Basel: Birkhäuser.

- Pfiffner, O. A., Lehner, P., Heitzmann, P., Müller, S., & Steck, A. (1997). *Deep structure of the Alps: Results of NRP20*. Basel: Birkhäuser Verlag.
- Philippe, Y., Deville, E., & Mascle, A. (1998). Thin-skinned inversion tectonics at oblique basin margins: example of the western Vercors and Chartreuse Subalpine massifs (SE France). *Geological Society, London, Special Publications*, 134, 239–262.
- Philipot, P. (1988). *Déformation et éclogitisation progressives d'une croûte océanique subductée: Le Monviso, Alpes occidentales*. Contraintes cinématiques durant la collision alpine (269 pp). Ph.D. thesis Université de Montpellier. Documents et Travaux Centre Géologique et Géophysique de Montpellier Numéro 19.
- Philipot, P. (1990). Opposite vergence of nappes and crustal extension in the French-Italian Western Alps. *Tectonics*, 9, 1143–1164.
- Piana, F. (2000). Structural Setting of Western Monferrato (Alps–Apennines Junction Zone, NW Italy). *Tectonics*, 19(5), 943–960.
- Piazza, A., Artoni, A., & Ogata, K. (2016). The Epiligurian wedge-top succession in the Enza Valley (Northern Apennines): evidence of a syn-depositional transpressive system. *Swiss Journal of Geosciences*, 109, 17–36.
- Picotti, V., & Pazzaglia, F. J. (2008). A new active tectonic model for the construction of the Northern Apennines mountain front near Bologna (Italy). *Journal of Geophysical Research*, 113, B08412. doi:10.1029/2007JB005307.
- Pieri, M. & Groppi, G. (1981). *Subsurface geological structure of the Po Plain, Italy*. Consiglio Nazionale di Ricerca, Progetto Finalizzato Geodinamica, 414, 13 pp.
- Platt, J. P., Behrmann, J. H., Cunningham, P. C., Dewey, J. F., Helman, M., Parish, M. G., et al. (1989). Kinematics of the Alpine arc and the motion history of Adria. *Nature*, 337, 158–161.
- Pleuger, J., Nagel, T.J., Walter, J.M., Jansen, E. & Froitzheim, N. (2008). On the role and importance of orogen-parallel and –perpendicular extension, transcurrent shearing, and backthrusting in the Monte Rosa nappe and the Southern Steep Belt of the Alps (Penninic zone, Switzerland and Italy). In S. Siegesmund, B. Fügenschuh & N. Froitzheim (Eds.), *Tectonic aspects of the Alpine-Dinaride-Carpathian System* (pp. 251–280). Geological Society London, Special Publications 298.
- Pleuger, J., Roller, S., Walter, J. M., Jansen, E., & Froitzheim, J. (2007). Structural evolution of the contact between two Penninic nappes (Zermatt-Saas zone and Combin zone, Western Alps) and implications for the exhumation mechanism and palaeogeography. *International Journal of Earth Sciences*, 96, 229–252.
- Pognante, U. (1989). Lawsonite, blueschist and eclogite formation in the southern Sesia Zone (western Alps, Italy). *European Journal Mineralogy*, 1, 89–104.
- Principi, G., Bortolotti, V., Chiari, M., Cortesogno, L., Gaggero, L., Marcucci, M., et al. (2004). The pre-orogenic volcano-sedimentary covers of the Western Tethys Oceanic basin: A review. *Ofioliti*, 29, 177–211.
- Principi, G., & Treves, B. (1984). Il sistema corso-appenninico come prisma d'accrescimento. Riflessi sul problema generale del limite Alpi-Appennini. *Memorie della Società Geologica Italiana*, 28, 549–576.
- Ramsay, J. G. (1980). Shear zone geometry: A review. *Journal of Structural Geology*, 2, 83–99.
- Regis, D., Rubato, D., Darling, J., Cenki-Tok, B., Zucali, M., & Engi, M. (2014). Multiple metamorphic stages within an eclogite-facies terrane (Sesia Zone, Western Alps) revealed by Th-U-Pb Petrochronology. *Journal of Petrology*, 55, 1429–1456.
- Ricou, L.E. & Siddans A.W.B. (1986). Collision tectonics in the western Alps. In M.P. Coward et al. (Eds.), *Collision tectonics* (pp. 229–244). Geological Society London Special Publication, 19.
- Ritz, J. F. (1992). Tectonique récente et sismotectonique des Alpes du Sud. *Quaternaire*, 3, 111–124.
- Rosenbaum, G., & Lister, G. S. (2004). Neogene and Quaternary rollback evolution of the Tyrrhenian Sea, the Apennines and the Sicilian Maghrebides. *Tectonics*, 23, TC1013. doi:10.1029/2003TC001518.
- Rosenberg, C. L., & Kissling, E. (2013). Three-dimensional insight into Central-Alpine collision: Lower-plate or upper-plate indentation? *Geology*, 41, 1219–1222. doi:10.1130/G34584.1.
- Rossetti, F., Faccenna, C., Jolivet, L., Funicello, R., Goffé, B., Tecce, F., et al. (2001). Structural signature and exhumation P-T-t path of the Gorgona island blueschist sequence (Tuscan Archipelago, Italy). *Ofioliti*, 26, 175–186.
- Rossetti, F., Faccenna, C., Jolivet, L., Goffé, B., & Funicello, R. (2002). Structural signature and exhumation P-T-t paths of the blueschist units exposed in the interior of the Northern Apennine chain, tectonic implications. *Bollettino Società Geologica Italiana Volume Speciale*, 1, 829–842.
- Roure, F., Bergerat, F., Damotte, Mugnier J.-L. & Polino R. (1996). The ECORS-CROP Alpine seismic traverse. *Mémoires Société Géologique de France*, 170, 113 pp.
- Roure, F., Polino, R., & Nicolich, R. (1990). Early Neogene deformation beneath the Po plain: Constraints on post-collisional Alpine evolution, in Deep Structure of the Alps. *Mémoires de la Société Géologique de France*, 156, 309–321.
- Rubato, D., & Hermann, J. (2001). Exhumation as fast as subduction? *Geology*, 29(1), 3–6.
- Scharf, A., Handy, M. R., Favaro, S., Schmid, S. M., & Bertrand, A. (2013). Modes of orogen-parallel stretching and extensional exhumation in response to microplate indentation and roll-back subduction (Tauern Window, Eastern Alps). *International Journal of Earth Sciences*, 102, 1627–1654.
- Schenk, V. (1990). The exposed crustal cross section of southern Calabria, Italy: Structure and evolution of a segment of Hercynian crust. In M. H. Salisbury & D. M. Fountain (Eds.), *Exposed cross-sections of the continental crust* (pp. 21–42). Netherlands: Kluwer Academic Publishers.
- Schmid, S.M., Aebli, H.R., Heller, F. & Zingg, A. (1989). The role of the Periadriatic Line in the tectonic evolution of the Alps. In M.P. Coward et al. (Eds.), *Alpine tectonics* (pp. 153–171). Geological Society London Special Publication 45.
- Schmid, S. M., Fügenschuh, B., Kissling, E., & Schuster, R. (2004). Tectonic map and overall architecture of the Alpine orogen. *Eclologiae Geologicae Helveticae*, 97, 93–117.
- Schmid, S. M., & Kissling, E. (2000). The arc of the Western Alps in the light of geophysical data on deep crustal structure. *Tectonics*, 19, 62–85.
- Schmid, S. M., Pfiffner, O. A., Froitzheim, N., Schönborn, G., & Kissling, E. (1996). Geophysical-geological transect and tectonic evolution of the Swiss-Italian Alps. *Tectonics*, 15, 1036–1064.
- Schmid, S. M., Zingg, A., & Handy, M. (1987). The kinematics of movements along the Insubric Line and the emplacement of the Ivrea Zone. *Tectonophysics*, 135, 47–66.
- Schönborn, G. (1992). Alpine tectonics and kinematic models of the central Southern Alps. *Memorie di Scienze Geologiche Padova*, 44, 229–393.
- Schumacher, M. E. (1997). Geological interpretation of the seismic profiles through the Southern Alps (lines S1-S7 and C3-south). In A. O. Pfiffner et al. (Eds.), *Deep structure of the Swiss Alps: Results from NRP 20* (pp. 101–114). Basel: Birkhäuser.
- Schumacher, M. E., & Laubscher, H. (1996). 3D crustal architecture of the Alps–Apennines join—a new view on seismic data. *Tectonophysics*, 260, 349–363.
- Schumacher, M. E., Schönborn, G., Bernoulli, D., & Laubscher, H. P. (1997). Rifting and collision in the Southern Alps. In A. O. Pfiffner, et al. (Eds.), *Deep Structure of the Swiss Alps:*

- Results From NRP 20 (pp. 186–204). Basel, Switzerland: Birkhäuser.
- Schwartz, S., Lardeaux, J. M., Tricart, P., Guillot, S., & Labrin, E. (2007). Diachronous exhumation of HP–LT metamorphic rocks from south-western Alps: evidence from fission-track analysis. *Terra Nova*, *19*, 133–140.
- Séranne, M. (1999). The Gulf of Lion continental margin (NW Mediterranean) revisited by IBS: An overview. *Geological Society, London, Special Publications*, *156*, 15–36.
- Serva, L. (Ed.) (2005). Gravity map of Italy and surrounding seas. Agenzia per la Protezione dell’Ambiente e per i Servizi Tecnici. Rome <http://www.isprambiente.gov.it>.
- Seton, M., Müller, R. D., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., et al. (2012). Global continental and ocean basin reconstructions since 200 Ma. *Earth-Science Reviews*, *113*, 212–270.
- Simon-Labric, T., Rolland, Y., Dumont, T., Heymes, T., Authemayou, C., Corsini, M., et al. (2009).  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of Penninic Front tectonic displacement (W Alps) during the Lower Oligocene (31–34 Ma). *Terra Nova*, *21*, 127–136.
- Skora, S., Mahlen, N. J., Johnson, C. M., Baumgartner, L. P., Lapen, T. J., Beard, B. L., et al. (2015). Evidence for protracted prograde metamorphism followed by rapid exhumation of the Zermatt-Saas Fee ophiolite. *Journal of Metamorphic Geology*, *33*, 711–734.
- Sommaruga, A. (1997). Geology of the central Jura and the Molasse basin: new insight into an evaporite-based foreland fold and thrust belt. *Mémoires Société Sciences Naturelles Neuchâtel*, *12*, 1–145.
- Sommaruga, A., Eichenberger, U. & Mariller, F. (2012). Seismic Atlas of the Swiss Molasse Basin. *Beiträge zur Geologie der Schweiz—Geophysik (text and 24 enclosures)*. Federal Office of Topography swisstopo.
- Spada, M., Bianchi, I., Kissling, E., Piana, Agostinetti N., & Wiemer, S. (2013). Combining controlled-source seismology and receiver function information to derive 3-D Moho topography for Italy. *Geophysical Journal International*, *194*, 1050–1068.
- Spakman, W., & Wortel, M. J. R. (2004). Tomographic view on western Mediterranean geodynamics. In W. Cavazza et al. (Eds.), *The TRANSMED Atlas, the mediterranean region from crust to mantle* (pp. 31–52). Berlin: Springer-Verlag.
- Steck, A., Della Torre, F., Keller, F., Pfeifer, H. R., Hunziker, J., & Masson, H. (2013). Tectonics of the Lepontine Alps: Ductile thrusting and folding in the deepest tectonic levels of the Central Alps. *Swiss Journal Geosciences*, *106*, 427–450.
- Steck, A., Epard, J.-L., Escher, A., Lehner, P., Marchant, R. & Masson, H. (1997). Geological interpretation of the seismic profiles through Western Switzerland: Rawil (W1), Val d’Anniviers (W2), Mattertal (W3), Zmutt-Zermatt-Findelen (W4) and Val de Bagnes (W5). In: Pfiffner A.O. et al. (Eds.): *Deep structure of the Swiss Alps: Results from NRP 20* (pp. 123–137). Basel: Birkhäuser.
- Steck, A., & Hunziker, J. C. (1994). The tertiary structural and thermal evolution of the Central Alps—compressional and extensional structures in an orogenic belt. *Tectonophysics*, *238*, 229–254.
- Steck, A., Masson, H., & Robyr, M. (2015). Tectonics of the Monte Rosa and surrounding nappes (Switzerland and Italy): Tertiary phases of subduction, thrusting and folding in the Pennine Alps. *Swiss Journal of Geosciences*, *108*, 3–34.
- Stipp, M., Stünitz, H., Heilbronner, R. & Schmid, S.M. (2002). Dynamic recrystallization of quartz: correlation between natural and experimental conditions. In De Meer et al. (Eds.), *Deformation mechanisms, rheology and tectonics: Current status and future perspectives* (pp. 171–190). Geological Society London Special Publication, 200.
- Strating, H. (1994). Extensional faulting in an intraoceanic subduction complex—working hypothesis for the Paleogene of the Alps–Apennine system. *Tectonophysics*, *238*, 255–273.
- Thouvenot, F., Senechal, G., Truffert, C., & Guellec, S. (1996). Comparison between two techniques of line-drawing migration (ray tracing and common tangent method). *Mémoires de la Société Géologique de France*, *170*, 53–59.
- Tricart, P. (1986). Le chevauchement de la zone briançonnaise au Sud-Est du Pelvoux: clé des rapports zone externe—zones internes dans les Alpes occidentales. *Bulletin de la Société géologique de France*, *11*, 233–244.
- Trullenque, G. (2005). *Tectonic and microfabric studies along the Penninic Front between Pelvoux and Argentera massifs (Western Alps)* (293 pp). University of Basel, Ph.D. thesis. Pdf available at <http://edoc.unibas.ch/353/>.
- Trullenque, G., Kunze, K., Heilbronner, R., Stünitz, H., & Schmid, S. M. (2006). Microfabrics of calcite ultramylonites as records of coaxial and non-coaxial deformation kinematics: Examples from the Rocher de l’Yret shear zone (Western Alps). *Tectonophysics*, *424*, 69–97.
- Van der Voo, R. (2004). Presidential Address: Paleomagnetism, oroclines, and growth of the continental crust. *Geological Society of America Today*, *14*(12), 4–9.
- van Hinsbergen, D. J. J., & Schmid, S. M. (2012). Map view restoration of Aegean–West Anatolian accretion and extension since the Eocene. *Tectonics*, *31*, TC5005. doi:10.1029/2012TC003132.
- van Hinsbergen, D. J. J., Vissers, R. L. M., & Spakman, W. (2014a). Origin and consequences of western Mediterranean subduction, rollback, and slab segmentation. *Tectonics*, *33*, 393–419.
- van Hinsbergen, D. J. J., Nebsink, M., Langereis, C. G., Maffione, M., Spalluto, L., Tropeano, M., & Sabato, L. (2014b). Did Adria rotate relative to Africa? *Solid Earth*, *5*, 611–629.
- van Wamel, W. A. (1987). On the tectonics of the Ligurian Apennines (northern Italy). *Tectonophysics*, *142*, 87–98.
- Vialon, P., Rochette, P. & Ménard, G. (1989). Indentation and rotation in the western Alpine arc. In M.P. Coward, D. Dietrich & R.G. Park (Eds.), *Alpine tectonics* (pp. 329–338). Geological Society of London Special Publications, 45.
- Villa, I. M., Bucher, S., Bousquet, R., Kleinhanns, I. C., & Schmid, S. M. (2014). Dating polygenic metamorphic assemblages along a transect across the Western Alps. *Journal of Petrology*, *55*, 803–830.
- Vitale Brovarone, A., & Herwartz, D. (2012). Timing of HP metamorphism in the Schistes Lustrés of Alpine Corsica: New Lu–Hf garnet and lawsonite ages. *Lithos*, *172–173*, 175–191.
- Vitale, S., & Ciarcia, S. (2013). Tectono-stratigraphic and kinematic evolution of the southern Apennines/Calabria–Peloritani Terrane system (Italy). *Tectonophysics*, *583*, 164–182.
- von Blanckenburg, F. (1992). Combined high-precision chronometry and geochemical tracing using accessory minerals: Applied to the Central-Alpine Bergell intrusion (central Europe). *Chemical Geology*, *100*, 19–40.
- Wagner, M., Kissling, E., & Husen, S. (2012). Combining controlled-source seismology and local earthquake tomography to derive a 3-D crustal model of the western Alpine region. *Geophysical Journal International*, *191*, 789–802.
- Walcott, C. R., & White, S. H. (1998). Constraints on the kinematics of post-orogenic extension imposed by stretching lineations in the Aegean region. *Tectonophysics*, *298*, 155–175.
- Weber, S., Sandmann, S., Miladinova, I., Fonseca, R. O. C., Froitzheim, N., Münker, C., et al. (2015). Dating the initiation of Piedmont-Liguria Ocean subduction: Lu–Hf garnet chronometry of eclogites from the Theodul Glacier Unit (Zermatt-Saas Zone, Switzerland). *Swiss Journal of Geosciences*, *108*, 183–199.

- Weissert, H. J., & Bemoulli, D. (1985). A transform margin in the Mesozoic Tethys: Evidence from the Swiss Alps. *Geologische Rundschau*, *71*, 665–679.
- Worthington, J. R., Hacker, B. R., & Zandt, G. (2013). Distinguishing eclogite from peridotite: EBSD-based calculations of seismic velocities. *Geophysical Journal International*, *193*, 489–505. doi:10.1093/gji/ggt004.
- Ye, S., Ansorge, J., Kissling, E., & Mueller, S. (1995). Crustal structure beneath the eastern Swiss Alps derived from seismic refraction data. *Tectonophysics*, *242*, 199–221.
- Zhao, L., Paul, A., Guillot, S., Solarino, S., Malusà, M. G., Zheng, T., et al. (2015). First seismic evidence for continental subduction beneath the Western Alps. *Geology*, *43*, 815–818.
- Zingg, A., Handy, M. R., Hunziker, J. C., & Schmid, S. M. (1990). Tectonometamorphic history of the Ivrea Zone and its relationship to the evolution of the Southern Alps. *Tectonophysics*, *182*, 169–192.